

Stellar winds of hot stars


Jiří Krtička

Masaryk University, Brno, Czech Republic



Observation of hot stars


- shells in the surroundings of hot stars



nebula close to the star WR 124 (HST)

Observation of hot stars


- the interstellar medium around hot stars



open cluster NGC 3603 (HST)

Observation of hot stars


- P Cyg line profiles in UV



HD 163758 (HST)

Observation of hot stars


- X-ray emission



X-ray spectrum θ^1 Ori C
(CHANDRA, Schulz et al. 2003)

Observation of hot stars


- H α emission line



α Cam, 2m telescope in Ondřejov (Kubát 2003)

How to explain the observations?

- nebulae



How to explain the observations?

- nebulae: circumstellar envelope around hot stars

How to explain the observations?

- nebulae: circumstellar envelope around hot stars
- influence on the interstellar medium




How to explain the observations?

- nebulae: circumstellar envelope around hot stars
- influence on the interstellar medium: envelope is expanding

How to explain the observations?

- nebulae: circumstellar envelope around hot stars
- influence on the interstellar medium: envelope is expanding
- P Cyg line profiles




How to explain the observations?

- nebulae: circumstellar envelope around hot stars
- influence on the interstellar medium: envelope is expanding
- P Cyg line profiles: supersonic outflow from hot stars: **wind**

How to explain the observations?

- nebulae: circumstellar envelope around hot stars
- influence on the interstellar medium: envelope is expanding
- P Cyg line profiles: supersonic outflow from hot stars: **wind**
- X-ray emission



How to explain the observations?

- nebulae: circumstellar envelope around hot stars
- influence on the interstellar medium: envelope is expanding
- P Cyg line profiles: supersonic outflow from hot stars: **wind**
- X-ray emission: shocks in the wind

How to explain the observations?

- nebulae: circumstellar envelope around hot stars
- influence on the interstellar medium: envelope is expanding
- P Cyg line profiles: supersonic outflow from hot stars: **wind**
- X-ray emission: shocks in the wind
- H α emission line

How to explain the observations?

- nebulae: circumstellar envelope around hot stars
- influence on the interstellar medium: envelope is expanding
- P Cyg line profiles: supersonic outflow from hot stars: **wind**
- X-ray emission: shocks in the wind
- $H\alpha$ emission line: recombination

How to explain the observations?

- nebulae: circumstellar envelope around hot stars
 - influence on the interstellar medium: envelope is expanding
 - P Cyg line profiles: supersonic outflow from hot stars: **wind**
 - X-ray emission: shocks in the wind
 - $H\alpha$ emission line: recombination
- ⇒ quantitative study of the wind

Hot star wind theory

- why is the wind blowing from hot stars?
- what are the main wind parameters (mass-loss rate, velocity)?
- how to predict the wind line profiles?
- how the wind influences the stellar evolution and the circumstellar environment?

Why is the wind blowing?

- some force accelerates the material from the stellar atmosphere to the circumstellar environment

Why is the wind blowing?

- hot stars are luminous: radiative force?

Why is the wind blowing?

- hot stars are luminous: radiative force?

$$f_{\text{rad}} = \frac{1}{c} \int_0^{\infty} \chi(r, \nu) F(r, \nu) d\nu$$

- spherically symmetric case
- $\chi(r, \nu)$ absorption coefficient
- $F(r, \nu)$ radiative flux

Why is the wind blowing?

- hot stars are luminous: radiative force?

$$f_{\text{rad}} = \frac{1}{c} \int_0^{\infty} \chi(r, \nu) F(r, \nu) d\nu$$

- radiative force due to the light scattering on free electrons

$$\chi(r, \nu) = \sigma_{\text{Th}} n_e(r)$$

- σ_{Th} Thomson scattering cross-section
- $n_e(r)$ electron density

Why is the wind blowing?

- hot stars are luminous: radiative force?

$$f_{\text{rad}} = \frac{1}{c} \int_0^{\infty} \chi(r, \nu) F(r, \nu) d\nu$$

- radiative force due to the light scattering on free electrons

$$f_{\text{rad}} = \frac{\sigma_{\text{Th}} n_e(r) L}{4\pi r^2 c}$$

$$\text{where } L = 4\pi r^2 \int_0^{\infty} F(r, \nu) d\nu$$

Why is the wind blowing?

- hot stars are luminous: radiative force?

$$f_{\text{rad}} = \frac{1}{c} \int_0^{\infty} \chi(r, \nu) F(r, \nu) d\nu$$

- radiative force due to the light scattering on free electrons

$$f_{\text{rad}} = \frac{\sigma_{\text{Th}} n_e(r) L}{4\pi r^2 c}$$

- comparison with the gravity force

$$f_{\text{grav}} = \frac{\rho(r) GM}{r^2}$$

Why is the wind blowing?

- hot stars are luminous: radiative force?

$$f_{\text{rad}} = \frac{1}{c} \int_0^{\infty} \chi(r, \nu) F(r, \nu) d\nu$$

- radiative force due to the light scattering on free electrons

$$f_{\text{rad}} = \frac{\sigma_{\text{T}} n_{\text{e}}(r) L}{4\pi r^2 c}$$

- comparison with the gravity force

$$\Gamma \equiv \frac{f_{\text{rad}}}{f_{\text{grav}}} = \frac{\sigma_{\text{T}} \frac{n_{\text{e}}(r)}{\rho(r)} L}{4\pi c G M}$$

Why is the wind blowing?

- hot stars are luminous: radiative force?

$$f_{\text{rad}} = \frac{1}{c} \int_0^{\infty} \chi(r, \nu) F(r, \nu) d\nu$$

- radiative force due to the light scattering on free electrons

$$f_{\text{rad}} = \frac{\sigma_{\text{Th}} n_e(r) L}{4\pi r^2 c}$$

- comparison with the gravity force

$$\Gamma \approx 10^{-5} \left(\frac{L}{1 L_{\odot}} \right) \left(\frac{M}{1 M_{\odot}} \right)^{-1}$$

Why is the wind blowing?

- hot stars are luminous: radiative force?

$$f_{\text{rad}} = \frac{1}{c} \int_0^{\infty} \chi(r, \nu) F(r, \nu) d\nu$$

- radiative force due to the light scattering on free electrons

$$f_{\text{rad}} = \frac{\sigma_{\text{Th}} n_e(r) L}{4\pi r^2 c}$$

- comparison with the gravity force
- example: α Cam, $L = 6.2 \times 10^5 L_{\odot}$,
 $M = 43 M_{\odot}$, $\Gamma \approx 0.1$

Why is the wind blowing?

- hot stars are luminous: radiative force?

$$f_{\text{rad}} = \frac{1}{c} \int_0^{\infty} \chi(r, \nu) F(r, \nu) d\nu$$

- radiative force due to the light scattering on free electrons

$$f_{\text{rad}} = \frac{\sigma_{\text{Th}} n_e(r) L}{4\pi r^2 c}$$

- comparison with the gravity force
- ⇒ radiative force due to the light scattering on free electrons is important, but it never (?) exceeds the gravity force

Why is the wind blowing?

- hot stars are luminous: radiative force?

$$f_{\text{rad}} = \frac{1}{c} \int_0^{\infty} \chi(r, \nu) F(r, \nu) d\nu$$

- radiative force due to the line transitions

$$\chi(r, \nu) = \frac{\pi e^2}{m_e c} \sum_{\text{lines}} \varphi_{ij}(\nu) g_i f_{ij} \left(\frac{n_i(r)}{g_i} - \frac{n_j(r)}{g_j} \right)$$

- $\varphi_{ij}(\nu)$ line profile, $\int_0^{\infty} \varphi_{ij}(\nu) = 1$
- f_{ij} oscillator strength
- $n_i(r)$, $n_j(r)$ level occupation number, g_i , g_j statistical weights

Why is the wind blowing?

- hot stars are luminous: radiative force?

$$f_{\text{rad}} = \frac{1}{c} \int_0^{\infty} \chi(r, \nu) F(r, \nu) d\nu$$

- radiative force due to the line transitions

$$f_{\text{line}} = \frac{\pi e^2}{m_e c^2} \int_0^{\infty} \sum_{\text{line}} g_i f_{ij} \left(\frac{n_i(r)}{g_i} - \frac{n_j(r)}{g_j} \right) \varphi_{ij}(\nu) F(r, \nu) d\nu$$

Why is the wind blowing?

- hot stars are luminous: radiative force?

$$f_{\text{rad}} = \frac{1}{c} \int_0^{\infty} \chi(r, \nu) F(r, \nu) d\nu$$

- radiative force due to the line transitions

$$f_{\text{line}} = \frac{\pi e^2}{m_e c^2} \int_0^{\infty} \sum_{\text{line}} g_i f_{ij} \left(\frac{n_i(r)}{g_i} - \frac{n_j(r)}{g_j} \right) \varphi_{ij}(\nu) F(r, \nu) d\nu$$

- problem: influence of lines on $F(r, \nu)$?

Why is the wind blowing?

- hot stars are luminous: radiative force?

$$f_{\text{rad}} = \frac{1}{c} \int_0^{\infty} \chi(r, \nu) F(r, \nu) d\nu$$

- radiative force due to the line transitions

$$f_{\text{line}} = \frac{\pi e^2}{m_e c^2} \int_0^{\infty} \sum_{\text{line}} g_i f_{ij} \left(\frac{n_i(r)}{g_i} - \frac{n_j(r)}{g_j} \right) \varphi_{ij}(\nu) F(r, \nu) d\nu$$

- problem: influence of lines on $F(r, \nu)$?
- **crude** solution: $F(r, \nu)$ constant for frequencies corresponding to a given line, $\nu \approx \nu_{ij}$

Why is the wind blowing?

- hot stars are luminous: radiative force?

$$f_{\text{rad}} = \frac{1}{c} \int_0^{\infty} \chi(r, \nu) F(r, \nu) d\nu$$

- radiative force due to the line transitions
 - maximum force

$$f_{\text{lines}}^{\text{max}} = \frac{\pi e^2}{m_e c^2} \sum_{\text{lines}} g_i f_{ij} \left(\frac{n_i(r)}{g_i} - \frac{n_j(r)}{g_j} \right) F(r, \nu_{ij})$$

- ν_{ij} is the line center frequency

Why is the wind blowing?

- hot stars are luminous: radiative force?

$$f_{\text{rad}} = \frac{1}{c} \int_0^{\infty} \chi(r, \nu) F(r, \nu) d\nu$$

- radiative force due to the line transitions
 - maximum force: comparison with gravity

$$\frac{f_{\text{line}}^{\text{max}}}{f_{\text{grav}}} = \frac{L e^2}{4 m_e \rho G M c^2} \sum_{\text{line}} f_{ij} n_i(r) \frac{L_\nu(\nu_{ij})}{L}$$

- neglect of $n_j(r) \ll n_i(r)$
- $L_\nu(\nu_{ij}) = 4\pi r^2 F(r, \nu_{ij})$

Why is the wind blowing?

- hot stars are luminous: radiative force?

$$f_{\text{rad}} = \frac{1}{c} \int_0^{\infty} \chi(r, \nu) F(r, \nu) d\nu$$

- radiative force due to the line transitions
 - maximum force: comparison with gravity

$$\frac{f_{\text{lines}}^{\text{max}}}{f_{\text{grav}}} = \Gamma \sum_{\text{lines}} \frac{\sigma_{ij}}{\sigma_{\text{Th}}} \frac{n_i}{n_e} \frac{\nu_{ij} L_{\nu}(\nu_{ij})}{L}$$

$$\sigma_{ij} = \frac{\pi e^2 f_{ij}}{\nu_{ij} m_e c}$$

Why is the wind blowing?

- hot stars are luminous: radiative force?

$$f_{\text{rad}} = \frac{1}{c} \int_0^{\infty} \chi(r, \nu) F(r, \nu) d\nu$$

- radiative force due to the line transitions
 - maximum force: comparison with gravity

$$\frac{f_{\text{lines}}^{\text{max}}}{f_{\text{grav}}} = \Gamma \sum_{\text{lines}} \frac{\sigma_{ij}}{\sigma_{\text{Th}}} \frac{n_i}{n_e} \frac{\nu_{ij} L_{\nu}(\nu_{ij})}{L}$$

- hydrogen: mostly ionised in the stellar envelopes $\Rightarrow n_i/n_e$ very small \Rightarrow negligible contribution to radiative force

Why is the wind blowing?

- hot stars are luminous: radiative force?

$$f_{\text{rad}} = \frac{1}{c} \int_0^{\infty} \chi(r, \nu) F(r, \nu) d\nu$$

- radiative force due to the line transitions
 - maximum force: comparison with gravity

$$\frac{f_{\text{lines}}^{\text{max}}}{f_{\text{grav}}} = \Gamma \sum_{\text{lines}} \frac{\sigma_{ij}}{\sigma_{\text{Th}}} \frac{n_i}{n_e} \frac{\nu_{ij} L_{\nu}(\nu_{ij})}{L}$$

- neutral helium: n_i/n_e very small \Rightarrow negligible contribution to radiative force

Why is the wind blowing?

- hot stars are luminous: radiative force?

$$f_{\text{rad}} = \frac{1}{c} \int_0^{\infty} \chi(r, \nu) F(r, \nu) d\nu$$

- radiative force due to the line transitions
 - maximum force: which elements?

$$\frac{f_{\text{lines}}^{\text{max}}}{f_{\text{grav}}} = \Gamma \sum_{\text{lines}} \frac{\sigma_{ij}}{\sigma_{\text{Th}}} \frac{n_i}{n_e} \frac{\nu_{ij} L_{\nu}(\nu_{ij})}{L}$$

- ionised helium: nonnegligible contribution to the radiative force

Why is the wind blowing?

- hot stars are luminous: radiative force?

$$f_{\text{rad}} = \frac{1}{c} \int_0^{\infty} \chi(r, \nu) F(r, \nu) d\nu$$

- radiative force due to the line transitions
 - maximum force: which elements?

$$\frac{f_{\text{lines}}^{\text{max}}}{f_{\text{grav}}} = \Gamma \sum_{\text{lines}} \frac{\sigma_{ij}}{\sigma_{\text{Th}}} \frac{n_i}{n_e} \frac{\nu_{ij} L_{\nu}(\nu_{ij})}{L}$$

- heavier elements (iron, carbon, nitrogen, oxygen, ...): large number of lines,
 $\sigma_{ij}/\sigma_{\text{Th}} \approx 10^7 \Rightarrow f_{\text{line}}^{\text{max}}/f_{\text{grav}}$ up to 10^3

Why is the wind blowing?

- hot stars are luminous: radiative force?

$$f_{\text{rad}} = \frac{1}{c} \int_0^{\infty} \chi(r, \nu) F(r, \nu) d\nu$$

- radiative force due to the line transitions
 - maximum force: which elements?

$$\frac{f_{\text{lines}}^{\text{max}}}{f_{\text{grav}}} = \Gamma \sum_{\text{lines}} \frac{\sigma_{ij}}{\sigma_{\text{Th}}} \frac{n_i}{n_e} \frac{\nu_{ij} L_{\nu}(\nu_{ij})}{L}$$

- ⇒ radiative force may be larger than gravity
(for many O stars $f_{\text{lines}}^{\text{max}} / f_{\text{grav}} \approx 2000$,
Abbott 1982, Gayley 1995)
- ⇒ **stellar wind**

Radiative force?

- speculations of Kepler, Newton

Radiative force?

- predicted by James Clerk Maxwell (1873) in the book *A Treatise on Electricity and Magnetism*



Radiative force?

- predicted by James Clerk Maxwell (1873)
- experimentally tested by Pyotr Nikolaevich Lebedev (1901), main problem: heating

Radiative force?

- predicted by James Clerk Maxwell (1873)
- experimentally tested by Pyotr Nikolaevich Lebedev (1901), main problem: heating
- why do we not observe the effects of the radiation pressure in a „normal world“?

Radiative force?

- predicted by James Clerk Maxwell (1873)
- experimentally tested by Pyotr Nikolaevich Lebedev (1901), main problem: heating
- why do we not observe the effects of the radiation pressure in a „normal world“?
 - classical particle: $E_p = \frac{1}{2}mv^2$, $p_p = \frac{2E}{v}$

Radiative force?

- predicted by James Clerk Maxwell (1873)
- experimentally tested by Pyotr Nikolaevich Lebedev (1901), main problem: heating
- why do we not observe the effects of the radiation pressure in a „normal world“?
 - classical particle: $E_p = \frac{1}{2}mv^2$, $p_p = \frac{2E}{v}$
 - photon: $E_\nu = h\nu$, $p_\nu = \frac{E}{c}$

Radiative force?

- predicted by James Clerk Maxwell (1873)
 - experimentally tested by Pyotr Nikolaevich Lebedev (1901), main problem: heating
 - why do we not observe the effects of the radiation pressure in a „normal world“?
 - classical particle: $E_p = \frac{1}{2}mv^2$, $p_p = \frac{2E}{v}$
 - photon: $E_\nu = h\nu$, $p_\nu = \frac{E}{c}$
- ⇒ for $E_p = E_\nu$ the momentum ratio is

$$\frac{p_\nu}{p_p} \approx \frac{v}{c}$$

Radiative force?

- predicted by James Clerk Maxwell (1873)
- experimentally tested by Pyotr Nikolaevich Lebedev (1901), main problem: heating
- why do we not observe the effects of the radiation pressure in a „normal world“?
 - particle with thermal energy $E_p \approx kT$

$$\frac{p_\nu}{p_p} \approx \frac{h\nu}{c\sqrt{mkT}} \approx 0.001 \left(\frac{\nu}{10^{15} \text{ s}^{-1}} \right) \left(\frac{T}{100 \text{ K}} \right)^{-1/2}$$

- two possibilities:

Radiative force?

- predicted by James Clerk Maxwell (1873)
- experimentally tested by Pyotr Nikolaevich Lebedev (1901), main problem: heating
- why do we not observe the effects of the radiation pressure in a „normal world“?
 - particle with thermal energy $E_p \approx kT$

$$\frac{p_\nu}{p_p} \approx \frac{h\nu}{c\sqrt{mkT}} \approx 0.001 \left(\frac{\nu}{10^{15} \text{ s}^{-1}} \right) \left(\frac{T}{100 \text{ K}} \right)^{-1/2}$$

- two possibilities:
 - large $\nu \Rightarrow$ X-rays, Compton effect

Radiative force?

- predicted by James Clerk Maxwell (1873)
- experimentally tested by Pyotr Nikolaevich Lebedev (1901), main problem: heating
- why do we not observe the effects of the radiation pressure in a „normal world“?
 - particle with thermal energy $E_p \approx kT$

$$\frac{p_\nu}{p_p} \approx \frac{h\nu}{c\sqrt{mkT}} \approx 0.001 \left(\frac{\nu}{10^{15} \text{ s}^{-1}} \right) \left(\frac{T}{100 \text{ K}} \right)^{-1/2}$$

- two possibilities:
 - large $\nu \Rightarrow$ X-rays, Compton effect
 - minimise heating (as did Lebedev)

Radiative force?

- predicted by James Clerk Maxwell (1873)
- experimentally tested by Pyotr Nikolaevich Lebedev (1901), main problem: heating
- why do we not observe the effects of the radiation pressure in a „normal world“?
 - how to minimise heating?

Radiative force?

- predicted by James Clerk Maxwell (1873)
- experimentally tested by Pyotr Nikolaevich Lebedev (1901), main problem: heating
- why do we not observe the effects of the radiation pressure in a „normal world“?
 - how to minimise heating?
 - cooling: emission of photon with the same energy as the absorbed one

Radiative force?

- predicted by James Clerk Maxwell (1873)
- experimentally tested by Pyotr Nikolaevich Lebedev (1901), main problem: heating
- why do we not observe the effects of the radiation pressure in a „normal world“?
 - how to minimise heating?
 - cooling: emission of photon with the same energy as the absorbed one
 - line absorption followed by emission
 - Thomson scattering


Radiative force?

- predicted by James Clerk Maxwell (1873)
- experimentally tested by Pyotr Nikolaevich Lebedev (1901), main problem: heating
- why do we not observe the effects of the radiation pressure in a „normal world“?
 - how to minimise heating?
 - cooling: emission of photon with the same energy as the absorbed one
 - line absorption followed by emission
 - Thomson scattering
 - both processes important in hot star winds

The Sobolev approximation


- the main problem: the line opacity (lines may be optically thick)
- ⇒ necessary to solve the radiative transfer equation

The Sobolev approximation




The Sobolev approximation

frequency ↑




velocity ↑



radius →


the Doppler effect in the wind

The Sobolev approximation




the Doppler effect in the wind

The Sobolev approximation




- $\Delta\nu_D$ is the Doppler width of the line

The Sobolev approximation




- $L_S \equiv \frac{v_{th}}{\frac{dv}{dr}} = c \frac{\Delta \nu_D}{\nu_{ij}} \frac{1}{\frac{dv}{dr}}$ is the Sobolev length

The Sobolev approximation




- structure does not significantly vary over $L_S \Rightarrow$ simplification of the calculation of f^{rad} possible

The Sobolev approximation



- opacity nonnegligible only over $L_S \Rightarrow$ solution of RTE in the „gray“ zone only

The Sobolev approximation



- $$H \equiv \frac{\rho}{\left(\frac{d\rho}{dr}\right)} \approx \frac{v}{\left(\frac{dv}{dr}\right)} \gg \frac{v_{th}}{\left(\frac{dv}{dr}\right)} \equiv L_S \quad (v \gg v_{th})$$

Our assumptions

- spherical symmetry

Our assumptions

- spherical symmetry
- stationary (time-independent) flow

The Sobolev line force I.

- the radiative transfer equation

$$\begin{aligned}\mu \frac{\partial}{\partial r} I(r, \mu, \nu) + \frac{1 - \mu^2}{r} \frac{\partial}{\partial \mu} I(r, \mu, \nu) &= \\ &= \eta(r, \mu, \nu) - \chi(r, \mu, \nu) I(r, \mu, \nu)\end{aligned}$$

- frame of static observer
- stationarity, spherical symmetry
- μ is frequency, $\mu = \cos \theta$
- $I(r, \mu, \nu)$ is specific intensity
- $\chi(r, \mu, \nu)$ is absorption (extinction) coefficient
- $\eta(r, \mu, \nu)$ is emissivity (emission coefficient)

The Sobolev line force I.

- the radiative transfer equation

$$\begin{aligned}\mu \frac{\partial}{\partial r} I(r, \mu, \nu) + \frac{1 - \mu^2}{r} \frac{\partial}{\partial \mu} I(r, \mu, \nu) &= \\ &= \eta(r, \mu, \nu) - \chi(r, \mu, \nu) I(r, \mu, \nu)\end{aligned}$$

- problem: $\chi(r, \mu, \nu)$ and $\eta(r, \mu, \nu)$ depend on μ due to the Doppler effect

The Sobolev line force I.

- the radiative transfer equation

$$\begin{aligned}\mu \frac{\partial}{\partial r} I(r, \mu, \nu) + \frac{1 - \mu^2}{r} \frac{\partial}{\partial \mu} I(r, \mu, \nu) &= \\ &= \eta(r, \mu, \nu) - \chi(r, \mu, \nu) I(r, \mu, \nu)\end{aligned}$$

- problem: $\chi(r, \mu, \nu)$ and $\eta(r, \mu, \nu)$ depend on μ due to the Doppler effect
- solution: use comoving frame!

The Sobolev line force I.

- CMF radiative transfer equation

$$\begin{aligned} \mu \frac{\partial}{\partial r} I(r, \mu, \nu) + \frac{1 - \mu^2}{r} \frac{\partial}{\partial \mu} I(r, \mu, \nu) - \\ \frac{\nu v(r)}{cr} \left(1 - \mu^2 + \frac{\mu^2 r}{v(r)} \frac{dv(r)}{dr} \right) \frac{\partial}{\partial \nu} I(r, \mu, \nu) = \\ = \eta(r, \nu) - \chi(r, \nu) I(r, \mu, \nu) \end{aligned}$$

- comoving frame (CMF) equation
- $v(r)$ is the fluid velocity
- $\chi(r, \nu)$ and $\eta(r, \nu)$ do depend on μ


The Sobolev line force I.

- CMF radiative transfer equation

$$\begin{aligned} \mu \frac{\partial}{\partial r} I(r, \mu, \nu) + \frac{1 - \mu^2}{r} \frac{\partial}{\partial \mu} I(r, \mu, \nu) - \\ \frac{\nu v(r)}{cr} \left(1 - \mu^2 + \frac{\mu^2 r}{v(r)} \frac{dv(r)}{dr} \right) \frac{\partial}{\partial \nu} I(r, \mu, \nu) = \\ = \eta(r, \nu) - \chi(r, \nu) I(r, \mu, \nu) \end{aligned}$$

- neglected aberration, advection (unimportant for $v \ll c$, e.g., Korčáková & Kubát 2003)
- neglect of the transformation of $I(r, \mu, \nu)$ between individual inertial frames

Intermezzo: the interpretation



- in CMF: continuous redshift of a given photon

The Sobolev line force II.

- the Sobolev transfer equation (Castor 2004)

$$\begin{aligned} & \mu \frac{\partial}{\partial r} I(r, \mu, \nu) + \frac{1 - \mu^2}{r} \frac{\partial}{\partial \mu} I(r, \mu, \nu) - \\ & - \frac{\nu v(r)}{cr} \left(1 - \mu^2 + \frac{\mu^2 r}{v(r)} \frac{dv(r)}{dr} \right) \frac{\partial}{\partial \nu} I(r, \mu, \nu) = \\ & = \eta(r, \nu) - \chi(r, \nu) I(r, \mu, \nu) \end{aligned}$$

The Sobolev line force II.

- the Sobolev transfer equation (Castor 2004)

$$\begin{aligned} & \mu \frac{\partial}{\partial r} I(r, \mu, \nu) + \frac{1 - \mu^2}{r} \frac{\partial}{\partial \mu} I(r, \mu, \nu) - \\ & - \frac{\nu v(r)}{cr} \left(1 - \mu^2 + \frac{\mu^2 r}{v(r)} \frac{dv(r)}{dr} \right) \frac{\partial}{\partial \nu} I(r, \mu, \nu) = \\ & = \eta(r, \nu) - \chi(r, \nu) I(r, \mu, \nu) \end{aligned}$$

- possible when $\frac{\nu v(r)}{cr} \frac{\partial}{\partial \nu} I(r, \mu, \nu) \gg \frac{\partial}{\partial r} I(r, \mu, \nu)$

- dimensional arguments:

- $\frac{\partial}{\partial r} I(r, \mu, \nu) \sim \frac{I(r, \mu, \nu)}{r},$

- $\frac{\partial}{\partial \nu} I(r, \mu, \nu) \sim \frac{I(r, \mu, \nu)}{\Delta \nu},$

$\Delta \nu = \nu \frac{v_{\text{th}}}{c}$ is the line Doppler width

The Sobolev line force II.

- the Sobolev transfer equation (Castor 2004)

$$\begin{aligned} & \mu \frac{\partial}{\partial r} I(r, \mu, \nu) + \frac{1 - \mu^2}{r} \frac{\partial}{\partial \mu} I(r, \mu, \nu) - \\ & - \frac{\nu v(r)}{cr} \left(1 - \mu^2 + \frac{\mu^2 r}{v(r)} \frac{dv(r)}{dr} \right) \frac{\partial}{\partial \nu} I(r, \mu, \nu) = \\ & = \eta(r, \nu) - \chi(r, \nu) I(r, \mu, \nu) \end{aligned}$$

- possible when $v(r) \gg v_{\text{th}}$

The Sobolev line force III.

- solution of the transfer equation for **one** line

$$\begin{aligned} -\frac{\nu v(r)}{cr} \left(1 - \mu^2 + \frac{\mu^2 r}{v(r)} \frac{dv(r)}{dr} \right) \frac{\partial}{\partial \nu} I(r, \mu, \nu) = \\ = \eta(r, \nu) - \chi(r, \nu) I(r, \mu, \nu) \end{aligned}$$

The Sobolev line force III.

- solution of the transfer equation for **one** line

$$\begin{aligned} -\frac{\nu v(r)}{cr} \left(1 - \mu^2 + \frac{\mu^2 r}{v(r)} \frac{dv(r)}{dr} \right) \frac{\partial}{\partial \nu} I(r, \mu, \nu) &= \\ &= \eta(r, \nu) - \chi(r, \nu) I(r, \mu, \nu) \end{aligned}$$

- line absorption and emission coefficients are

$$\chi(r, \nu) = \frac{\pi e^2}{m_e c} \varphi_{ij}(\nu) g_i f_{ij} \left(\frac{n_i(r)}{g_i} - \frac{n_j(r)}{g_j} \right)$$

$$\eta(r, \nu) = \frac{2h\nu^3}{c^2} \frac{\pi e^2}{m_e c} \varphi_{ij}(\nu) g_i f_{ij} \frac{n_j(r)}{g_j}$$

The Sobolev line force III.

- solution of the transfer equation for **one** line

$$\begin{aligned} -\frac{\nu v(r)}{cr} \left(1 - \mu^2 + \frac{\mu^2 r}{v(r)} \frac{dv(r)}{dr} \right) \frac{\partial}{\partial \nu} I(r, \mu, \nu) &= \\ &= \eta(r, \nu) - \chi(r, \nu) I(r, \mu, \nu) \end{aligned}$$

- the line opacity and emissivity are

$$\chi(r, \nu) = \chi_L(r) \varphi_{ij}(\nu)$$

$$\eta(r, \nu) = \chi_L(r) S_L(r) \varphi_{ij}(\nu)$$

$$\text{where } \chi_L(r) = \frac{\pi e^2}{m_e c} g_i f_{ij} \left(\frac{n_i(r)}{g_i} - \frac{n_j(r)}{g_j} \right)$$

The Sobolev line force III.

- solution of the transfer equation for **one** line

$$\begin{aligned} -\frac{\nu v(r)}{cr} \left(1 - \mu^2 + \frac{\mu^2 r}{v(r)} \frac{dv(r)}{dr} \right) \frac{\partial}{\partial \nu} I(r, \mu, \nu) = \\ = \chi_L(r) \varphi_{ij}(\nu) (S_L(r) - I(r, \mu, \nu)) \end{aligned}$$

The Sobolev line force III.

- solution of the transfer equation for **one** line

$$\begin{aligned} -\frac{\nu v(r)}{cr} \left(1 - \mu^2 + \frac{\mu^2 r}{v(r)} \frac{dv(r)}{dr} \right) \frac{\partial}{\partial \nu} I(r, \mu, \nu) &= \\ &= \chi_L(r) \varphi_{ij}(\nu) (S_L(r) - I(r, \mu, \nu)) \end{aligned}$$

- introduce a new variable

$$y = \int_{\nu}^{\infty} d\nu' \varphi_{ij}(\nu')$$

- where
 - $y = 0$: the incoming side of the line
 - $y = 1$: the outgoing side of the line

The Sobolev line force III.

- solution of the transfer equation for **one** line

$$\begin{aligned} \frac{\nu v(r)}{cr} \left(1 - \mu^2 + \frac{\mu^2 r}{v(r)} \frac{dv(r)}{dr} \right) \frac{\partial}{\partial y} I(r, \mu, y) = \\ = \chi_L(r) (S_L(r) - I(r, \mu, y)) \end{aligned}$$

The Sobolev line force III.

- solution of the transfer equation for **one** line

$$\begin{aligned} \frac{\nu v(r)}{cr} \left(1 - \mu^2 + \frac{\mu^2 r}{v(r)} \frac{dv(r)}{dr} \right) \frac{\partial}{\partial y} I(r, \mu, y) = \\ = \chi_L(r) (S_L(r) - I(r, \mu, y)) \end{aligned}$$

- assumptions:
 - variables do not significantly vary with r within the „resonance zone“

$$\Rightarrow \text{fixed } r, \frac{\partial}{\partial y} \rightarrow \frac{d}{dy}$$

- $\nu \rightarrow \nu_0$

\Rightarrow integration possible

The Sobolev line force III.

- solution of the transfer equation for **one** line


$$I(y) = I_c(\mu) \exp[-\tau(\mu)y] + S_L \{1 - \exp[-\tau(\mu)y]\}$$

- where
 - the Sobolev optical depth is

$$\tau(\mu) = \frac{\chi_L(r)cr}{\nu_0 v(r) \left(1 - \mu^2 + \frac{\mu^2 r}{v(r)} \frac{dv(r)}{dr}\right)}$$

- the boundary condition is $I(y = 0) = I_c(\mu)$

Intermezzo: the interpretation



- τ is given by the slope $\Rightarrow \tau \sim \left(\frac{dv}{dr}\right)^{-1}$

The Sobolev line force IV.

- the radiative force (the radial component; force per unit of volume)

$$f_{\text{rad}} = \frac{1}{c} \int_0^{\infty} \chi(r, \nu) F(r, \nu) d\nu$$

The Sobolev line force IV.

- the radiative force (the radial component; force per unit of volume)

$$f_{\text{rad}} = \frac{1}{c} \int_0^{\infty} d\nu \chi(r, \nu) \oint d\Omega \mu I(r, \mu, \nu)$$

The Sobolev line force IV.

- the radiative force (the radial component; force per unit of volume)

$$f_{\text{rad}} = \frac{2\pi}{c} \int_0^{\infty} d\nu \chi_L(r) \varphi_{ij}(\nu) \int_{-1}^1 d\mu \mu I(r, \mu, \nu)$$

The Sobolev line force IV.

- the radiative force (the radial component; force per unit of volume)

$$f_{\text{rad}} = \frac{2\pi\chi_L(r)}{c} \int_0^1 dy \int_{-1}^1 d\mu \mu I(r, \mu, y)$$

The Sobolev line force IV.

- the radiative force (the radial component; force per unit of volume)

$$f_{\text{rad}} = \frac{2\pi\chi_L(r)}{c} \int_0^1 dy \times \int_{-1}^1 d\mu \mu \{ I_c(\mu) \exp[-\tau(\mu)y] + S_L \{1 - \exp[-\tau(\mu)y]\} \}$$

- where the Sobolev optical depth is

$$\tau(\mu) = \frac{\chi_L(r)cr}{\nu_0 v(r) \left(1 - \mu^2 + \frac{\mu^2 r}{v(r)} \frac{dv(r)}{dr} \right)}$$

- $\tau(\mu)$ is an even function of μ

The Sobolev line force IV.

- the radiative force (the radial component; force per unit of volume)

$$f_{\text{rad}} = \frac{2\pi\chi_L(r)}{c} \int_0^1 dy \int_{-1}^1 d\mu \mu I_c(\mu) \exp[-\tau(\mu)y]$$

- no net contribution of the emission to the radiative force (S_L is isotropic in the CMF)

The Sobolev line force IV.

- the radiative force (the radial component; force per unit of volume)

$$f_{\text{rad}} = \frac{2\pi\chi_L(r)}{c} \int_{-1}^1 d\mu \mu I_c(\mu) \frac{1 - \exp[-\tau(\mu)]}{\tau(\mu)}$$

- inserting

$$\tau(\mu) = \frac{\chi_L(r)cr}{\nu_0 v(r) \left(1 - \mu^2 + \frac{\mu^2 r}{v(r)} \frac{dv(r)}{dr} \right)}$$

The Sobolev line force IV.

- the radiative force (the radial component; force per unit of volume)

$$f_{\text{rad}} = \frac{2\pi\nu_0 v(r)}{rc^2} \int_{-1}^1 d\mu \mu I_c(\mu) [1 + \mu^2 \sigma(r)] \times \\ \times \left\{ 1 - \exp \left[-\frac{\chi_L(r) cr}{\nu_0 v(r) (1 + \mu^2 \sigma(r))} \right] \right\}$$

- where $\sigma(r) = \frac{r}{v(r)} \frac{dv(r)}{dr} - 1$
- Sobolev (1957), Castor (1974), Rybicki & Hummer (1978)

Optically thin lines

- optically thin line:

$$\frac{\chi_L(r)cr}{\nu_0 v(r) (1 + \mu^2 \sigma(r))} \ll 1$$

Optically thin lines

- optically thin line:

$$\frac{\chi_L(r)cr}{\nu_0 v(r) (1 + \mu^2 \sigma(r))} \ll 1$$

- the radiative force proportional to

$$f_{\text{rad}} \sim 1 - \exp \left[-\frac{\chi_L(r)cr}{\nu_0 v(r) (1 + \mu^2 \sigma(r))} \right]$$

Optically thin lines

- optically thin line:

$$\frac{\chi_L(r)cr}{\nu_0 v(r) (1 + \mu^2 \sigma(r))} \ll 1$$

- the radiative force proportional to

$$f_{\text{rad}} \sim 1 - \exp \left[-\frac{\chi_L(r)cr}{\nu_0 v(r) (1 + \mu^2 \sigma(r))} \right]$$
$$\approx \frac{\chi_L(r)cr}{\nu_0 v(r) (1 + \mu^2 \sigma(r))}$$

Optically thin lines

$$f_{\text{rad}} = \frac{2\pi}{c} \int_{-1}^1 d\mu \mu I_c(\mu) \chi_L(r)$$

Optically thin lines

$$f_{\text{rad}} = \frac{1}{c} \chi_{\text{L}}(r) F(r)$$

Optically thin lines

$$f_{\text{rad}} = \frac{1}{c} \chi_{\text{L}}(r) F(r)$$

- optically thin radiative force proportional to the radiative flux $F(r)$
- optically thin radiative force proportional to the normalised line opacity $\chi_{\text{L}}(r)$ (or to the density)
- the same result as for the static medium

Optically thick lines

- optically thick line:

$$\frac{\chi_L(r)cr}{\nu_0 v(r) (1 + \mu^2 \sigma(r))} \gg 1$$

Optically thick lines

- optically thick line:

$$\frac{\chi_L(r)cr}{\nu_0 v(r) (1 + \mu^2 \sigma(r))} \gg 1$$

- the radiative force proportional to

$$f_{\text{rad}} \sim 1 - \exp \left[-\frac{\chi_L(r)cr}{\nu_0 v(r) (1 + \mu^2 \sigma(r))} \right]$$

Optically thick lines

- optically thick line:

$$\frac{\chi_L(r)cr}{\nu_0 v(r) (1 + \mu^2 \sigma(r))} \gg 1$$

- the radiative force proportional to

$$f_{\text{rad}} \sim 1 - \exp \left[-\frac{\chi_L(r)cr}{\nu_0 v(r) (1 + \mu^2 \sigma(r))} \right]$$
$$\approx 1$$

Optically thick lines

$$f_{\text{rad}} = \frac{2\pi\nu_0 v(r)}{rc^2} \int_{-1}^1 d\mu \mu I_c(\mu) [1 + \mu^2 \sigma(r)]$$

Optically thick lines

$$f_{\text{rad}} = \frac{2\pi\nu_0 v(r)}{rc^2} \int_{-1}^1 d\mu \mu I_c(\mu) [1 + \mu^2 \sigma(r)]$$

- neglect of the limb darkening:

$$I_c(\mu) = \begin{cases} I_c = \text{const.}, & \mu \geq \mu_*, \\ 0, & \mu < \mu_* \end{cases},$$

where $\mu_* = \sqrt{1 - \frac{R_*^2}{r^2}}$

Optically thick lines

$$f_{\text{rad}} = \frac{2\pi\nu_0 v(r)}{rc^2} \int_{\mu_*}^1 d\mu \mu l_c [1 + \mu^2 \sigma(r)]$$

Optically thick lines

$$f_{\text{rad}} = \frac{\nu_0 v(r) F(r)}{rc^2} \left[1 + \sigma(r) \left(1 - \frac{1}{2} \frac{R_*^2}{r^2} \right) \right]$$

where $F = 2\pi \int_{\mu_*}^1 d\mu \mu l_c = \pi \frac{R_*^2}{r^2} l_c$

Optically thick lines

$$f_{\text{rad}} = \frac{\nu_0 v(r) F(r)}{rc^2} \left[1 + \sigma(r) \left(1 - \frac{1}{2} \frac{R_*^2}{r^2} \right) \right]$$

- large distance from the star: $r \gg R_*$

Optically thick lines

$$f_{\text{rad}} = \frac{\nu_0 v(r) F(r)}{rc^2} \left[1 + \sigma(r) \left(1 - \frac{1}{2} \frac{R_*^2}{r^2} \right) \right]$$

- large distance from the star: $r \gg R_*$

$$f_{\text{rad}} \approx \frac{\nu_0 F(r)}{c^2} \frac{dv(r)}{dr}$$

Optically thick lines

$$f_{\text{rad}} = \frac{\nu_0 v(r) F(r)}{rc^2} \left[1 + \sigma(r) \left(1 - \frac{1}{2} \frac{R_*^2}{r^2} \right) \right]$$

- large distance from the star: $r \gg R_*$

$$f_{\text{rad}} \approx \frac{\nu_0 F(r)}{c^2} \frac{dv(r)}{dr}$$

- optically thick radiative force proportional to the radiative flux $F(r)$
- optically thick radiative force proportional to $\frac{dv}{dr}$
- optically thick radiative force does not depend on the level populations or the density

Wind driven by thick lines

- continuity and momentum equation of isothermal spherically symmetric wind

$$\frac{\partial \rho}{\partial t} + \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 \rho v) = 0$$

$$\frac{\partial v}{\partial t} + \rho v \frac{\partial v}{\partial r} = -a^2 \frac{\partial \rho}{\partial r} + f_{\text{rad}} - \frac{\rho G M (1 - \Gamma)}{r^2}$$

- ρ, v are the wind density and velocity
- a is the sound speed

Wind driven by thick lines

- continuity and momentum equation of isothermal spherically symmetric wind

$$\frac{1}{r^2} \frac{d}{dr} (r^2 \rho v) = 0$$

$$\rho v \frac{dv}{dr} = -a^2 \frac{d\rho}{dr} + f_{\text{rad}} - \frac{\rho G M (1 - \Gamma)}{r^2}$$

- assumption: stationary flow

Wind driven by thick lines

- continuity equation

$$\frac{1}{r^2} \frac{d}{dr} (r^2 \rho v) = 0 \Rightarrow \dot{M} \equiv 4\pi r^2 \rho v = \text{const.}$$

- \dot{M} is the wind **mass-loss rate**

Wind driven by thick lines

- momentum equation

$$v \frac{dv}{dr} = \frac{f_{\text{rad}}}{\rho} - \frac{GM(1 - \Gamma)}{r^2}$$

- neglect of the gas-pressure term $a^2 \frac{d\rho}{dr} \ll f_{\text{rad}}$
(possible in the supersonic part of the wind)

Wind driven by thick lines

- momentum equation

$$v \frac{dv}{dr} = \frac{\nu_0 v(r) F(r)}{\rho r c^2} \left[1 + \sigma(r) \left(1 - \frac{1}{2} \frac{R_*^2}{r^2} \right) \right] - \frac{GM(1 - \Gamma)}{r^2}$$

- inclusion of the expression for the optically thick line force
- $F(r) = \frac{L_\nu}{4\pi r^2}$, where L_ν is the monochromatic stellar luminosity (constant)
- $\sigma(r) = \frac{r}{v} \frac{dv}{dr} - 1$

Wind driven by thick lines

- momentum equation

$$\left[v - \frac{\nu_0 L_\nu}{4\pi r^2 \rho c^2} \left(1 - \frac{1}{2} \frac{R_*^2}{r^2} \right) \right] \frac{dv}{dr} = \frac{\nu_0 v(r) L_\nu}{8\pi \rho c^2 r^3} - \frac{GM(1 - \Gamma)}{r^2}$$

Wind driven by thick lines

- momentum equation

$$\left[v - \frac{\nu_0 L_\nu}{4\pi r^2 \rho c^2} \left(1 - \frac{1}{2} \frac{R_*^2}{r^2} \right) \right] \frac{dv}{dr} = \frac{\nu_0 v(r) L_\nu}{8\pi \rho c^2 r^3} - \frac{GM(1 - \Gamma)}{r^2}$$

- has a **critical point**

$$v - \frac{\nu_0 L_\nu}{4\pi r^2 \rho c^2} \left(1 - \frac{1}{2} \frac{R_*^2}{r^2} \right) = 0$$

Wind driven by thick lines

- momentum equation

$$\left[v - \frac{\nu_0 L_\nu}{4\pi r^2 \rho c^2} \left(1 - \frac{1}{2} \frac{R_*^2}{r^2} \right) \right] \frac{dv}{dr} = \frac{\nu_0 v(r) L_\nu}{8\pi \rho c^2 r^3} - \frac{GM(1 - \Gamma)}{r^2}$$

- has a **critical point**

$$v - \frac{\nu_0 L_\nu}{4\pi r^2 \rho c^2} \left(1 - \frac{1}{2} \frac{R_*^2}{r^2} \right) = 0$$

- neglect of $\frac{R_*}{r}$ term:

$$\dot{M} \equiv 4\pi r^2 \rho v(r) = \frac{\nu_0 L_\nu}{c^2}$$

Wind driven by thick lines

- momentum equation

$$\left[v - \frac{\nu_0 L_\nu}{4\pi r^2 \rho c^2} \left(1 - \frac{1}{2} \frac{R_*^2}{r^2} \right) \right] \frac{dv}{dr} = \frac{\nu_0 v(r) L_\nu}{8\pi \rho c^2 r^3} - \frac{GM(1 - \Gamma)}{r^2}$$

- has a **critical point**


$$v - \frac{\nu_0 L_\nu}{4\pi r^2 \rho c^2} \left(1 - \frac{1}{2} \frac{R_*^2}{r^2} \right) = 0$$

- neglect of $\frac{R_*}{r}$ term:

$$\dot{M} \equiv 4\pi r^2 \rho v(r) = \frac{\nu_0 L_\nu}{c^2} \approx \frac{L}{c^2}$$

⇒ mass-loss rate due to one optically thick line
approximatively equal to the „photon mass-loss
rate“ (L is stellar luminosity)

Example: α Cam



Example: α Cam

| | |
|------------------------------|----------|
| temperature T_{eff} | 30 900 K |
|------------------------------|----------|

| | |
|--------------|------------------|
| radius R_* | 27.6 R_{\odot} |
|--------------|------------------|

| | |
|----------|----------------|
| mass M | 43 M_{\odot} |
|----------|----------------|

(Lamers et al. 1995)

Example: α Cam

| | |
|------------------------------|------------------|
| temperature T_{eff} | 30 900 K |
| radius R_* | 27.6 R_{\odot} |
| mass M | 43 M_{\odot} |

- mass-loss rate due to one optically thick line
 $\dot{M} \approx L/c^2$

Example: α Cam

| | |
|------------------------------|------------------|
| temperature T_{eff} | 30 900 K |
| radius R_* | 27.6 R_{\odot} |
| mass M | 43 M_{\odot} |

- mass-loss rate due to one optically thick line
 $\dot{M} \approx L/c^2$
- mass-loss rate due to N_{thick} optically thick lines
 $\dot{M} \approx N_{\text{thick}} L/c^2$

Example: α Cam

| | |
|------------------------------|------------------|
| temperature T_{eff} | 30 900 K |
| radius R_* | 27.6 R_{\odot} |
| mass M | 43 M_{\odot} |

- mass-loss rate due to one optically thick line
 $\dot{M} \approx L/c^2$
- mass-loss rate due to N_{thick} optically thick lines
 $\dot{M} \approx N_{\text{thick}} L/c^2$
- NLTE calculations: $N_{\text{thick}} \approx 1000$

Example: α Cam

| | |
|------------------------------|------------------|
| temperature T_{eff} | 30 900 K |
| radius R_* | 27.6 R_{\odot} |
| mass M | 43 M_{\odot} |

- mass-loss rate due to one optically thick line
 $\dot{M} \approx L/c^2$
- mass-loss rate due to N_{thick} optically thick lines
 $\dot{M} \approx N_{\text{thick}} L/c^2$
- NLTE calculations: $N_{\text{thick}} \approx 1000$
- $L = 4\pi\sigma R_*^2 T_{\text{eff}}^4$, $L = 620\,000 L_{\odot}$

Example: α Cam

| | |
|------------------------------|------------------|
| temperature T_{eff} | 30 900 K |
| radius R_* | 27.6 R_{\odot} |
| mass M | 43 M_{\odot} |

- mass-loss rate due to one optically thick line
 $\dot{M} \approx L/c^2$
- mass-loss rate due to N_{thick} optically thick lines
 $\dot{M} \approx N_{\text{thick}} L/c^2$
- NLTE calculations: $N_{\text{thick}} \approx 1000$
- $L = 4\pi\sigma R_*^2 T_{\text{eff}}^4$, $L = 620\,000 L_{\odot}$
- $\dot{M} \approx 4 \times 10^{-5} M_{\odot} \text{ yr}^{-1}$, more precise estimate:
 $1.5 \times 10^{-6} M_{\odot} \text{ yr}^{-1}$ (Krtička & Kubát 2008)

CAK theory

- in reality the wind is driven by a mixture of optically thick and thin lines
 - optically thin line force

$$f_{\text{rad}} = \frac{1}{c} \chi_L(r) F(r)$$

- optically thick line force

$$f_{\text{rad}} = \frac{\nu_0 F(r)}{c^2} \frac{dv}{dr}$$

CAK theory

- in reality the wind is driven by a mixture of optically thick and thin lines
 - optically thin line force

$$f_{\text{rad}} = \frac{1}{c} \chi_L(r) F(r)$$

- optically thick line force

$$f_{\text{rad}} = \frac{\nu_0 F(r)}{c^2} \frac{dv}{dr}$$

- Sobolev optical depth $\tau_S = \frac{\chi_L(r)c}{\nu_0 \frac{dv}{dr}}$

$$f_{\text{rad}} = \frac{1}{c} \chi_L(r) F(r) (\tau_S^{-1})^\alpha$$

where $\alpha = 0$ (thin) or $\alpha = 1$ (thick)

CAK theory

- in reality the wind is driven by a mixture of optically thick and thin lines

$$\Rightarrow 0 < \alpha < 1$$

CAK theory

- in reality the wind is driven by a mixture of optically thick and thin lines
- the radiative force in the **CAK approximation** (Castor, Abbott & Klein 1975)

$$f_{\text{rad}} = k \frac{\sigma_{\text{Th}} n_e L}{4\pi r^2 c} \left(\frac{1}{\sigma_{\text{Th}} n_e v_{\text{th}}} \frac{dv}{dr} \right)^\alpha$$

- where
 - k, α are constants (force multipliers)
 - σ_{Th} is the Thomson scattering cross-section
 - n_e is the electron number density
 - v_{th} is hydrogen thermal speed (for $T = T_{\text{eff}}$)
- (Abbott 1982)

CAK theory

- in reality the wind is driven by a mixture of optically thick and thin lines
- the radiative force in the **CAK approximation** (Castor, Abbott & Klein 1975)

$$f_{\text{rad}} = k \frac{\sigma_{\text{Th}} n_e L}{4\pi r^2 c} \left(\frac{1}{\sigma_{\text{Th}} n_e v_{\text{th}}} \frac{dv}{dr} \right)^\alpha$$

- nondimensional parameters k and α describe the line-strength distribution function (CAK, Puls et al. 2000)
- in general NLTE calculations necessary to obtain k and α (Abbott 1982)

CAK theory

- momentum equation with CAK line force (neglecting the gas pressure term)

$$\rho v \frac{dv}{dr} = f_{\text{rad}} - \frac{\rho GM(1 - \Gamma)}{r^2}$$

CAK theory

- momentum equation with CAK line force (neglecting the gas pressure term)

$$\rho v \frac{dv}{dr} = k \frac{\sigma_{\text{Th}} n_e L}{4\pi r^2 c} \left(\frac{1}{\sigma_{\text{Th}} n_e v_{\text{th}}} \frac{dv}{dr} \right)^\alpha - \frac{\rho G M (1 - \Gamma)}{r^2}$$

CAK theory

- momentum equation with CAK line force (neglecting the gas pressure term)

$$r^2 v \frac{dv}{dr} = k \frac{\sigma_{\text{Th}} L n_e}{4\pi c \rho} \left(\frac{\rho}{n_e \sigma_{\text{Th}} \dot{M} v_{\text{th}}} \frac{4\pi r^2 v}{dr} \frac{dv}{dr} \right)^\alpha - GM(1 - \Gamma)$$

CAK theory

- momentum equation with CAK line force (neglecting the gas pressure term)

$$r^2 v \frac{dv}{dr} = k \frac{\sigma_{\text{Th}} L n_e}{4\pi c \rho} \left(\frac{\rho}{n_e \sigma_{\text{Th}} \dot{M} v_{\text{th}}} \frac{dv}{dr} \right)^\alpha - GM(1 - \Gamma)$$

- velocity in terms of the escape speed

$$w \equiv \frac{v^2}{v_{\text{esc}}^2}, \text{ where } v_{\text{esc}}^2 = \frac{2GM(1 - \Gamma)}{R_*}$$

- new radial variable

$$x \equiv 1 - \frac{R_*}{r}$$

CAK theory

- momentum equation with CAK line force (neglecting the gas pressure term)

$$1 + w' = C (w')^\alpha$$

- where

- $w' \equiv \frac{dw}{dx}$

- $C \equiv \frac{k\sigma_{\text{Th}}L}{4\pi cGM(1-\Gamma)} \frac{n_e}{\rho} \left(\frac{\rho}{n_e} \frac{4\pi GM(1-\Gamma)}{\sigma_{\text{Th}}\dot{M}v_{\text{th}}} \right)^\alpha$

- $\frac{\rho}{n_e} \approx m_{\text{H}}$

- algebraic equation

CAK theory

- momentum equation with CAK line force (neglecting the gas pressure term)


$$1 + w' = C (w')^\alpha$$

- different solutions for different values of C (or mass-loss rate \dot{M})

CAK theory

- momentum equation with CAK line force (neglecting the gas pressure term)


$$1 + w' = C (w')^\alpha$$



CAK theory

- momentum equation with CAK line force (neglecting the gas pressure term)

$$1 + w' = C (w')^\alpha$$




- large C (small \dot{M}): two solutions

CAK theory

- momentum equation with CAK line force (neglecting the gas pressure term)

$$1 + w' = C (w')^\alpha$$




- small C (large \dot{M}): no solution

CAK theory

- momentum equation with CAK line force (neglecting the gas pressure term)

$$1 + w' = C (w')^\alpha$$



- critical value of $C (\dot{M})$: one solution

CAK theory

- momentum equation with CAK line force (neglecting the gas pressure term)

$$1 + w' = C (w')^\alpha$$

- critical (CAK) solution for a specific value of \dot{M} : the only smooth solution of detailed momentum equation from the stellar surface to infinity
- CAK solution: the largest \dot{M} possible

CAK theory

- momentum equation with CAK line force (neglecting the gas pressure term)

$$1 + w' = C (w')^\alpha$$

- critical (CAK) solution for a specific value of \dot{M} : the only smooth solution of detailed momentum equation from the stellar surface to infinity
- ⇒ possible to derive the wind mass-loss rate and velocity profile

$$w'_c = \frac{\alpha}{1 - \alpha}$$

$$C_c = \frac{(1 - \alpha)^{\alpha-1}}{\alpha^\alpha}$$

CAK theory


$$w'_c = \frac{\alpha}{1 - \alpha}$$

$$\Rightarrow w = \frac{\alpha}{1 - \alpha} x \Rightarrow v = v_\infty \left(1 - \frac{R_*}{r}\right)^{1/2}$$

- where the terminal velocity

$$v_\infty = v_{\text{esc}} \sqrt{\frac{\alpha}{1 - \alpha}}$$

CAK theory



CAK theory

$$w'_c = \frac{\alpha}{1 - \alpha}$$

$$\Rightarrow w = \frac{\alpha}{1 - \alpha} x \Rightarrow v = v_\infty \left(1 - \frac{R_*}{r}\right)^{1/2}$$

- where the terminal velocity

$$v_\infty = v_{\text{esc}} \sqrt{\frac{\alpha}{1 - \alpha}}$$

- v_∞ scales with v_{esc} !

CAK theory

$$w'_c = \frac{\alpha}{1 - \alpha}$$

$$\Rightarrow w = \frac{\alpha}{1 - \alpha} x \Rightarrow v = v_\infty \left(1 - \frac{R_*}{r}\right)^{1/2}$$

- where the terminal velocity

$$v_\infty = v_{\text{esc}} \sqrt{\frac{\alpha}{1 - \alpha}}$$

- v_∞ scales with v_{esc} !
- as v_∞ of order of 100 km s^{-1} , hot star winds are strongly supersonic!

CAK theory

$$w'_c = \frac{\alpha}{1 - \alpha}$$

$$\Rightarrow w = \frac{\alpha}{1 - \alpha} x \Rightarrow v = v_\infty \left(1 - \frac{R_*}{r}\right)^{1/2}$$

- where the terminal velocity

$$v_\infty = v_{\text{esc}} \sqrt{\frac{\alpha}{1 - \alpha}}$$

- v_∞ scales with v_{esc} !
- example: α Cam, $v_{\text{esc}} = 620 \text{ km s}^{-1}$, $\alpha = 0.61$

CAK theory

$$w'_c = \frac{\alpha}{1 - \alpha}$$

$$\Rightarrow w = \frac{\alpha}{1 - \alpha} x \Rightarrow v = v_\infty \left(1 - \frac{R_*}{r}\right)^{1/2}$$

- where the terminal velocity

$$v_\infty = v_{\text{esc}} \sqrt{\frac{\alpha}{1 - \alpha}}$$

- v_∞ scales with v_{esc} !
- example: α Cam, $v_{\text{esc}} = 620 \text{ km s}^{-1}$, $\alpha = 0.61$
 \Rightarrow prediction: $v_\infty = 780 \text{ km s}^{-1}$

CAK theory

$$C_c = \frac{(1 - \alpha)^{\alpha-1}}{\alpha^\alpha}$$

$$\Rightarrow \dot{M} = \left[\frac{4\pi m_H GM(1 - \Gamma)}{\sigma_{Th}} \right]^{\frac{\alpha-1}{\alpha}} \frac{\alpha}{v_{th} (1 - \alpha)^{\frac{\alpha-1}{\alpha}}} \left(\frac{kL}{c} \right)^{\frac{1}{\alpha}}$$

CAK theory

$$C_c = \frac{(1 - \alpha)^{\alpha-1}}{\alpha^\alpha}$$

$$\Rightarrow \dot{M} = \left[\frac{4\pi m_H G M (1 - \Gamma)}{\sigma_{\text{Th}}} \right]^{\frac{\alpha-1}{\alpha}} \frac{\alpha}{v_{\text{th}} (1 - \alpha)^{\frac{\alpha-1}{\alpha}}} \left(\frac{kL}{c} \right)^{\frac{1}{\alpha}}$$

- example: α Cam: $\dot{M} \approx 9 \times 10^{-6} M_\odot \text{ yr}^{-1}$

Beyond the classical CAK theory

- inclusion of the dependence of k on the ionisation equilibrium – δ parameter (Abbott 1982)

Beyond the classical CAK theory

- inclusion of the dependence of k on the ionisation equilibrium – δ parameter (Abbott 1982)
- dropping of the radial streaming approximation (Pauldrach, Puls & Kudritzki 1986, Friend & Abbott 1986)

Beyond the classical CAK theory

- inclusion of the dependence of k on the ionisation equilibrium – δ parameter (Abbott 1982)
- dropping of the radial streaming approximation (Pauldrach, Puls & Kudritzki 1986, Friend & Abbott 1986)
- NLTE calculation of the level populations (Pauldrach 1987, Vink, de Koter & Lamers 2000, Gräfener & Hamann 2002, Krtićka & Kubát 2004)

Beyond the classical CAK theory

- inclusion of the dependence of k on the ionisation equilibrium – δ parameter (Abbott 1982)
- dropping of the radial streaming approximation (Pauldrach, Puls & Kudritzki 1986, Friend & Abbott 1986)
- NLTE calculation of the level populations (Pauldrach 1987, Vink, de Koter & Lamers 2000, Gräfener & Hamann 2002, Krtićka & Kubát 2004)
- dropping of the Sobolev approximation (Pauldrach et al. 1994, Gräfener & Hamann 2002)

Comparison with observations

- nice wind theory \Rightarrow compare it with observations!

Comparison with observations

- nice wind theory \Rightarrow compare it with observations!
- time for hot chocolate (observers will do the work for us)!



Comparison with observations

- nice wind theory \Rightarrow compare it with observations!
- time for hot chocolate (observers will do the work for us)!?

no coffee time yet...

Comparison with observations


- nice wind theory \Rightarrow compare it with observations!
 - time for hot chocolate (observers will do the work for us)!?
 - problem: it is not possible to „measure“ the wind parameters directly from observations
- \Rightarrow we have to work more to understand the wind spectral characteristics

Comparison with observations


- nice wind theory \Rightarrow compare it with observations!
 - time for hot chocolate (observers will do the work for us)!?
 - problem: it is not possible to „measure“ the wind parameters directly from observations
- \Rightarrow we have to work more to understand the wind spectral characteristics
- more theory, please!

Observations: H α line profiles


- H α emission line of α Cam




Observations: H α line profiles




Observations: H α line profiles



Observations: H α line profiles



Observations: H α line profiles



Observations: $H\alpha$ line profiles

- our assumption: $H\alpha$ line is optically thin

Observations: H α line profiles

- our assumption: H α line is optically thin
- number of H α photons emitted per unit of time

$$N_{\text{H}\alpha} \sim n_p n_e$$

- where
 - n_p is the number density of H $^+$
 - n_e is the number density of free electrons

Observations: H α line profiles

- our assumption: H α line is optically thin
- number of H α photons emitted per unit of time

$$N_{\text{H}\alpha} \sim n_p n_e$$

- as $n_p \sim \dot{M}$ and $n_e \sim \dot{M} \Rightarrow N_{\text{H}\alpha} \sim \dot{M}^2$

Observations: H α line profiles

- our assumption: H α line is optically thin
- number of H α photons emitted per unit of time

$$N_{\text{H}\alpha} \sim n_p n_e$$

- as $n_p \sim \dot{M}$ and $n_e \sim \dot{M} \Rightarrow N_{\text{H}\alpha} \sim \dot{M}^2$
- \Rightarrow possibility to derive \dot{M} using NLTE models

Observations: H α line profiles


- our assumption: H α line is optically thin
- number of H α photons emitted per unit of time

$$N_{\text{H}\alpha} \sim n_p n_e$$

- as $n_p \sim \dot{M}$ and $n_e \sim \dot{M} \Rightarrow N_{\text{H}\alpha} \sim \dot{M}^2$
- \Rightarrow possibility to derive \dot{M} using NLTE models
- example: α Cam
 - our estimate: $9 \times 10^{-6} M_{\odot} \text{ yr}^{-1}$
 - theoretical prediction: $1.4 \times 10^{-6} M_{\odot} \text{ yr}^{-1}$
(Krtićka & Kubát 2007)
 - H α line observation: $1.5 \times 10^{-6} M_{\odot} \text{ yr}^{-1}$
(Puls et al. 2006)

Observations: P Cyg lines I.


- IUE spectrum of α Cam




- saturated line profile of P Cyg type

Observations: P Cyg lines I.



- lines of the most abundant ion of a given element




Observations: P Cyg lines I.



Observations: P Cyg lines I.




Observations: P Cyg lines I.



Observations: P Cyg lines I.


- IUE spectrum of α Cam



- absorption in the wind between star and observer
- emission due to the wind around the star

Observations: P Cyg lines I.


- IUE spectrum of α Cam



- the absorption edge originates in the wind with the highest velocity in the direction of observer

Observations: P Cyg lines I.


- IUE spectrum of α Cam



- the absorption edge originates in the wind with the highest velocity in the direction of observer
- possibility to derive the terminal velocity v_∞


Observations: P Cyg lines I.

- IUE spectrum of α Cam



Observations: P Cyg lines I.

- IUE spectrum of α Cam




$$v_\infty = \frac{\Delta\lambda}{\lambda_0} c$$

- where λ_0 is the laboratory wavelength of a given line

Observations: P Cyg lines I.

- IUE spectrum of α Cam




$$v_\infty = \frac{\Delta\lambda}{\lambda_0} c$$

- α Cam: $\Delta\lambda = 7.9 \text{ \AA} \Rightarrow v_\infty = 1500 \text{ km s}^{-1}$
- our estimate: 780 km s^{-1}

Observations: P Cyg lines I.


- IUE spectrum of α Cam



- why is the absorption part saturated?

Observations: P Cyg lines I.

- IUE spectrum of α Cam




- why is the absorption part saturated?

$$I(y) = I_c(\mu) \exp[-\tau(\mu)y] + S_L \{1 - \exp[-\tau(\mu)y]\}$$

- the emergent intensity: $y \rightarrow 1$

Observations: P Cyg lines I.

- IUE spectrum of α Cam




- why is the absorption part saturated?

$$I = I_c(\mu) \exp[-\tau(\mu)] + S_L \{1 - \exp[-\tau(\mu)]\}$$

- optically thick lines $\tau \gg 1$ with $S_L \ll I_c \Rightarrow I \ll I_c$

Observations: P Cyg lines I.


- IUE spectrum of α Cam



- for saturated lines ($\tau \gg 1$) the absorption part of the P Cyg line profile does not depend on τ
 - ⇒ determination of v_∞ possible
 - ⇒ determination of \dot{M} impossible

Observations: P Cyg lines II.


- HST spectrum of HD 13268



- unsaturated line profile of P Cyg type

Observations: P Cyg lines II.

- HST spectrum of HD 13268




$$\frac{F}{F_c} \approx \exp[-\tau(\mu = 1)]$$

$$\tau(\mu = 1) = \frac{\chi_{LC}}{\nu_0} \left(\frac{dv}{dr} \right)^{-1}$$

Observations: P Cyg lines II.

- HST spectrum of HD 13268




$$\frac{F}{F_c} \approx \exp[-\tau(\mu = 1)]$$

$$\tau(\mu = 1) = \frac{\pi e^2}{m_e c} g_i f_{ij} \left(\frac{n_i(r)}{g_i} - \frac{n_j(r)}{g_j} \right) \frac{c}{\nu_0} \left(\frac{dv}{dr} \right)^{-1}$$

Observations: P Cyg lines II.

- HST spectrum of HD 13268




$$\frac{F}{F_c} \approx \exp[-\tau(\mu = 1)]$$

$$\tau(\mu = 1) = \frac{\pi e^2}{m_e c} \lambda_{ij} f_{ij} n_i(r) \left(\frac{dv}{dr} \right)^{-1}$$

Observations: P Cyg lines II.

- HST spectrum of HD 13268




$$\tau(\mu = 1) = \frac{\pi e^2}{m_e c} \lambda_{ij} f_{ij} \frac{q_{\text{CIV}} Z_C \dot{M}}{4\pi m_H v r^2} \left(\frac{dv}{dr} \right)^{-1}$$

- Z_C is the carbon number density relatively to H
- q_{CIV} is the ionisation fraction of CIV

Observations: P Cyg lines II.

- HST spectrum of HD 13268




$$\tau(\mu = 1) = \frac{\pi e^2}{m_e c} \lambda_{ij} f_{ij} \frac{Z_C}{4\pi m_H} \frac{1}{v_\infty^2 R_*} q_{CIV} \dot{M}$$

- our order-of-magnitude approximations:
 $v \rightarrow v_\infty$, $r \rightarrow R_*$, $dv/dr \rightarrow v_\infty/R_*$

Observations: P Cyg lines II.

- HST spectrum of HD 13268




$$\tau(\mu = 1) = \frac{\pi e^2}{m_e c} \lambda_{ij} f_{ij} \frac{Z_C}{4\pi m_H} \frac{1}{v_\infty^2 R_*} q_{\text{CIV}} \dot{M}$$

\Rightarrow from unsaturated wind line profiles possible to derive $q_{\text{CIV}} \dot{M}$

Observations: P Cyg lines II.

- HST spectrum of HD 13268




$$\tau(\mu = 1) = \frac{\pi e^2}{m_e c} \lambda_{ij} f_{ij} \frac{Z_C}{4\pi m_H} \frac{1}{v_\infty^2 R_*} q_{CIV} \dot{M}$$

- in our case $q_{CIV} \dot{M} = 4 \times 10^{-10} M_\odot \text{ yr}^{-1}$
- \dot{M} can be derived with a knowledge of q_{CIV}

Observation: X-ray emission

- X-ray spectrum θ^1 Ori C



(CHANDRA, Schulz et al. 2003)

Observation: X-ray emission

- X-ray emission of hot stars consists of numerous lines of highly excited elements (N VI, O VII, Fe XXIV, . . .)
- signature of a presence of gas with temperatures of the order 10^6 K
- X-ray emission originates in the wind
 - how?

Observation: X-ray emission

- problem:
 - the wind temperature is of the order of the stellar effective temperature – 10^4 K (as expected from the observed ionisation structure and as derived from NLTE models, e.g., Drew 1989)
 - how can such gas emit X-ray radiation with typical temperatures $\sim 10^6$ K?

Observation: X-ray emission

- problem:
 - the wind temperature is of the order of the stellar effective temperature – 10^4 K
 - how can such gas emit X-ray radiation with typical temperatures $\sim 10^6$ K?
- solution:
 - most of the wind material is „cool“ with temperatures of order of 10^4 K
 - only a very small fraction of the wind is very hot $\sim 10^6$ K
 - the „hot“ material quickly cools down (radiatively)

Observation: X-ray emission


- problem:
 - the wind temperature is of the order of the stellar effective temperature – 10^4 K
 - how can such gas emit X-ray radiation with typical temperatures $\sim 10^6$ K?
- solution:
 - most of the wind material is „cool“ with temperatures of order of 10^4 K
 - only a very small fraction of the wind is very hot $\sim 10^6$ K
 - the „hot“ material quickly cools down (radiatively)
- further problem: how is this possible?

How to create X-rays?

- hot stars have stellar wind with typical velocities $\approx 1000 \text{ km s}^{-1}$


How to create X-rays?

- hot stars have stellar wind with typical velocities $\approx 1000 \text{ km s}^{-1}$



How to create X-rays?


- hot stars have stellar wind with typical velocities $\approx 1000 \text{ km s}^{-1}$



$T = 2 \cdot 10^7 \text{ K}$


Can wind material collide?

- possible influence of the wind instabilities



Can wind material collide?

- possible influence of the wind instabilities



Wind instabilities I.

- main idea
 - the Sobolev approximation gives reliable prediction of wind structure
- ⇒ a sound basis for the study of instabilities

Wind instabilities I.

- time-dependent hydrodynamical equations

$$\frac{\partial \rho}{\partial t} + \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 \rho v) = 0$$

$$\rho \frac{\partial v}{\partial t} + \rho v \frac{\partial v}{\partial r} = -a^2 \frac{\partial \rho}{\partial r} + f_{\text{rad}} - \frac{\rho G M (1 - \Gamma)}{r^2}$$

- ρ, v are the wind density and velocity
- a is the sound speed

Wind instabilities I.

- time-dependent hydrodynamical equations

$$\frac{\partial \rho}{\partial t} + \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 \rho v) = 0$$

$$\rho \frac{\partial v}{\partial t} + \rho v \frac{\partial v}{\partial r} = -a^2 \frac{\partial \rho}{\partial r} + f_{\text{rad}} - \frac{\rho G M (1 - \Gamma)}{r^2}$$

- comoving fluid-frame + small perturbations of stationary solution
 - $\rho = \rho_0 + \delta\rho,$
 - $v = v_0 + \delta v, v_0 = 0$

Wind instabilities I.

- equations for perturbations $\delta\rho$, δv

$$\frac{\partial\delta\rho}{\partial t} + \rho_0 \frac{\partial\delta v}{\partial r} = 0$$

$$\rho_0 \frac{\partial\delta v}{\partial t} = -a^2 \frac{\partial\delta\rho}{\partial r} + \delta f_{\text{rad}}$$

- perturbation of the radiative force
 $\delta f_{\text{rad}} = \rho_0 g'_{\text{rad}} \delta v / \delta r$
- where $g'_{\text{rad}} \equiv \partial g_{\text{rad}} / \partial (dv/dr)$

Wind instabilities I.

- the wave equation

$$\frac{\partial^2 \delta v}{\partial t^2} = a^2 \frac{\partial^2 \delta v}{\partial r^2} + g'_{\text{rad}} \frac{\partial^2 \delta v}{\partial t \partial r}$$

Wind instabilities I.

- the wave equation

$$\frac{\partial^2 \delta v}{\partial t^2} = a^2 \frac{\partial^2 \delta v}{\partial r^2} + g'_{\text{rad}} \frac{\partial^2 \delta v}{\partial t \partial r}$$

- solution in the form $\delta v \sim \exp [i (\omega t - kr)]$

Wind instabilities I.

- the wave equation

$$\frac{\partial^2 \delta v}{\partial t^2} = a^2 \frac{\partial^2 \delta v}{\partial r^2} + g'_{\text{rad}} \frac{\partial^2 \delta v}{\partial t \partial r}$$

- the dispersion relation

$$\omega^2 + g'_{\text{rad}} \omega k - a^2 k^2 = 0$$

Wind instabilities I.

- the wave equation

$$\frac{\partial^2 \delta v}{\partial t^2} = a^2 \frac{\partial^2 \delta v}{\partial r^2} + g'_{\text{rad}} \frac{\partial^2 \delta v}{\partial t \partial r}$$

- the dispersion relation

$$\frac{\omega}{k} = -\frac{1}{2} g'_{\text{rad}} \pm \left(\frac{1}{4} g'^2_{\text{rad}} + a^2 \right)^{1/2}$$

Wind instabilities I.

- the wave equation

$$\frac{\partial^2 \delta v}{\partial t^2} = a^2 \frac{\partial^2 \delta v}{\partial r^2} + g'_{\text{rad}} \frac{\partial^2 \delta v}{\partial t \partial r}$$

- the dispersion relation

$$\frac{\omega}{k} = -\frac{1}{2} g'_{\text{rad}} \pm \left(\frac{1}{4} g'^2_{\text{rad}} + a^2 \right)^{1/2}$$

- zero radiative force

$$\frac{\omega}{k} = \pm a$$

- ordinary sound waves

Wind instabilities I.

- the wave equation

$$\frac{\partial^2 \delta v}{\partial t^2} = a^2 \frac{\partial^2 \delta v}{\partial r^2} + g'_{\text{rad}} \frac{\partial^2 \delta v}{\partial t \partial r}$$

- the dispersion relation

$$\frac{\omega}{k} = -\frac{1}{2} g'_{\text{rad}} \pm \left(\frac{1}{4} g'^2_{\text{rad}} + a^2 \right)^{1/2}$$

- general case
 - new type of waves – radiative-acoustic (Abbott) waves (Abbott 1980, Feldmeier et al. 2008)
 - downstream (+) and upstream (-) mode

Wind instabilities I.

- the wave equation

$$\frac{\partial^2 \delta v}{\partial t^2} = a^2 \frac{\partial^2 \delta v}{\partial r^2} + g'_{\text{rad}} \frac{\partial^2 \delta v}{\partial t \partial r}$$

- the dispersion relation

$$\frac{\omega}{k} = -\frac{1}{2} g'_{\text{rad}} \pm \left(\frac{1}{4} g'^2_{\text{rad}} + a^2 \right)^{1/2}$$

- **critical point:** radial wind velocity equals to the speed of (upstream) Abbott waves

$$v_c - \frac{1}{2} g'_{\text{rad}} - \left(\frac{1}{4} g'^2_{\text{rad}} + a^2 \right)^{1/2} = 0$$

Wind instabilities I.

- the wave equation

$$\frac{\partial^2 \delta v}{\partial t^2} = a^2 \frac{\partial^2 \delta v}{\partial r^2} + g'_{\text{rad}} \frac{\partial^2 \delta v}{\partial t \partial r}$$

- the dispersion relation

$$\frac{\omega}{k} = -\frac{1}{2} g'_{\text{rad}} \pm \left(\frac{1}{4} g'^2_{\text{rad}} + a^2 \right)^{1/2}$$

- **critical point**: radial wind velocity equals to the speed of (upstream) Abbott waves
- ⇒ no information can travel from the regions with $v > v_c$ towards the stellar surface (critical surface resembles the event horizon of a black hole, Feldmeier & Shlosman 2000)

Wind instabilities I.

- the wave equation

$$\frac{\partial^2 \delta v}{\partial t^2} = a^2 \frac{\partial^2 \delta v}{\partial r^2} + g'_{\text{rad}} \frac{\partial^2 \delta v}{\partial t \partial r}$$

- the dispersion relation

$$\frac{\omega}{k} = -\frac{1}{2} g'_{\text{rad}} \pm \left(\frac{1}{4} g'^2_{\text{rad}} + a^2 \right)^{1/2}$$

- **critical point**: radial wind velocity equals to the speed of (upstream) Abbott waves
- ⇒ no information can travel from the regions with $v > v_c$ towards the stellar surface
- ⇒ mass-loss rate is determined there

Wind instabilities I.

- the wave equation

$$\frac{\partial^2 \delta v}{\partial t^2} = a^2 \frac{\partial^2 \delta v}{\partial r^2} + g'_{\text{rad}} \frac{\partial^2 \delta v}{\partial t \partial r}$$

- the dispersion relation

$$\frac{\omega}{k} = -\frac{1}{2} g'_{\text{rad}} \pm \left(\frac{1}{4} g'^2_{\text{rad}} + a^2 \right)^{1/2}$$

⇒ no instability of hot-star winds!

Wind instabilities I.

- the wave equation

$$\frac{\partial^2 \delta v}{\partial t^2} = a^2 \frac{\partial^2 \delta v}{\partial r^2} + g'_{\text{rad}} \frac{\partial^2 \delta v}{\partial t \partial r}$$

- the dispersion relation

$$\frac{\omega}{k} = -\frac{1}{2} g'_{\text{rad}} \pm \left(\frac{1}{4} g'^2_{\text{rad}} + a^2 \right)^{1/2}$$

⇒ no instability of hot-star winds!

- hydrodynamical simulations
(Votruba et al. 2007)

Wind instabilities II.

- our stability analysis showed that the wind should be stable

Wind instabilities II.

- our stability analysis showed that the wind should be stable
- what causes the occurrence of X-rays?


Wind instabilities II.

- our stability analysis showed that the wind should be stable
- what causes the occurrence of X-rays?
- what is wrong with our stability analysis?

Wind instabilities II.


- our stability analysis showed that the wind should be stable
- what causes the occurrence of X-rays?
- what is wrong with our stability analysis?
- the Sobolev approximation is not valid for small (optically thin) perturbations!

Wind instabilities II.




- the radiative transfer in the comoving frame

Wind instabilities II.




- the absorption profile in the comoving frame

Wind instabilities II.



- the line force

Wind instabilities II.



- the line force after a small change of the velocity

Wind instabilities II.

- the radiative acceleration

$$g_{\text{rad}} = \frac{2\pi}{c\rho} \int_0^\infty d\nu \chi_L(r) \varphi_{ij}(\nu) \int_{-1}^1 d\mu \mu I(r, \mu, \nu)$$

Wind instabilities II.

- the radiative acceleration

$$g_{\text{rad}} = \frac{2\pi}{c\rho} \int_0^\infty d\nu \chi_L(r) \varphi_{ij}(\nu) \int_{-1}^1 d\mu \mu I(r, \mu, \nu)$$

- optically thin perturbation

$$\delta g_{\text{rad}} = \frac{2\pi}{c\rho} \int_0^\infty d\nu \chi_L(r) \delta\varphi_{ij}(\nu) \int_{-1}^1 d\mu \mu I(r, \mu, \nu)$$

Wind instabilities II.

- the radiative acceleration

$$g_{\text{rad}} = \frac{2\pi}{c\rho} \int_0^\infty d\nu \chi_L(r) \varphi_{ij}(\nu) \int_{-1}^1 d\mu \mu I(r, \mu, \nu)$$

- optically thin perturbation

$$\delta g_{\text{rad}} = \frac{2\pi}{c\rho} \int_0^\infty d\nu \chi_L(r) \delta \varphi_{ij}(\nu) \int_{-1}^1 d\mu \mu I(r, \mu, \nu)$$

$$\delta \varphi_{ij}(\nu) = \frac{d\varphi_{ij}(\nu)}{d\nu} \delta \nu = \frac{d\varphi_{ij}(\nu)}{d\nu} \nu_0 \frac{\delta v}{c}$$

Wind instabilities II.

- the radiative acceleration

$$g_{\text{rad}} = \frac{2\pi}{c\rho} \int_0^\infty d\nu \chi_L(r) \varphi_{ij}(\nu) \int_{-1}^1 d\mu \mu I(r, \mu, \nu)$$

- optically thin perturbation

$$\delta g_{\text{rad}} = \frac{2\pi}{c\rho} \int_0^\infty d\nu \chi_L(r) \delta \varphi_{ij}(\nu) \int_{-1}^1 d\mu \mu I(r, \mu, \nu)$$

$$\delta \varphi_{ij}(\nu) = \frac{d\varphi_{ij}(\nu)}{d\nu} \delta \nu = \frac{d\varphi_{ij}(\nu)}{d\nu} \nu_0 \frac{\delta v}{c}$$

$$\Rightarrow \delta g_{\text{rad}} = \Omega \delta v \quad (\Omega > 0)$$

Wind instabilities II.

- equations for perturbations $\delta\rho$, δv

$$\frac{\partial\delta\rho}{\partial t} + \rho_0 \frac{\partial\delta v}{\partial r} = 0$$

$$\rho_0 \frac{\partial\delta v}{\partial t} = -a^2 \frac{\partial\delta\rho}{\partial r} + \delta f_{\text{rad}}$$

Wind instabilities II.

- the wave equation

$$\frac{\partial^2 \delta v}{\partial t^2} = a^2 \frac{\partial^2 \delta v}{\partial r^2} + \Omega \frac{\partial \delta v}{\partial t}$$

Wind instabilities II.

- the wave equation

$$\frac{\partial^2 \delta v}{\partial t^2} = a^2 \frac{\partial^2 \delta v}{\partial r^2} + \Omega \frac{\partial \delta v}{\partial t}$$

- solution in the form $\delta v \sim \exp [i (\omega t - kr)]$

Wind instabilities II.

- the wave equation

$$\frac{\partial^2 \delta v}{\partial t^2} = a^2 \frac{\partial^2 \delta v}{\partial r^2} + \Omega \frac{\partial \delta v}{\partial t}$$

- the dispersion relation

$$\omega^2 + i\Omega\omega - a^2 k^2 = 0$$

Wind instabilities II.

- the wave equation

$$\frac{\partial^2 \delta v}{\partial t^2} = a^2 \frac{\partial^2 \delta v}{\partial r^2} + \Omega \frac{\partial \delta v}{\partial t}$$

- the dispersion relation

$$\omega = -\frac{1}{2}i\Omega \pm \left(-\frac{1}{4}\Omega^2 + a^2 k^2 \right)^{1/2}$$

Wind instabilities II.

- the wave equation

$$\frac{\partial^2 \delta v}{\partial t^2} = a^2 \frac{\partial^2 \delta v}{\partial r^2} + \Omega \frac{\partial \delta v}{\partial t}$$

- the dispersion relation

$$\omega = -\frac{1}{2}i\Omega \pm \left(-\frac{1}{4}\Omega^2 + a^2 k^2 \right)^{1/2}$$

- negligible gas pressure: $\Omega^2 \gg a^2 k^2$

Wind instabilities II.

- the wave equation

$$\frac{\partial^2 \delta v}{\partial t^2} = a^2 \frac{\partial^2 \delta v}{\partial r^2} + \Omega \frac{\partial \delta v}{\partial t}$$

- the dispersion relation (non-zero ω)

$$\omega = -i\Omega$$

Wind instabilities II.

- the wave equation

$$\frac{\partial^2 \delta v}{\partial t^2} = a^2 \frac{\partial^2 \delta v}{\partial r^2} + \Omega \frac{\partial \delta v}{\partial t}$$

- the dispersion relation (non-zero ω)

$$\omega = -i\Omega$$

- the wave amplitude varies as ($\Omega > 0$)

$$\delta v \sim \exp(i\omega t) = \exp(\Omega t)$$

Wind instabilities II.

- the wave equation

$$\frac{\partial^2 \delta v}{\partial t^2} = a^2 \frac{\partial^2 \delta v}{\partial r^2} + \Omega \frac{\partial \delta v}{\partial t}$$

- the dispersion relation (non-zero ω)

$$\omega = -i\Omega$$

- the wave amplitude varies as ($\Omega > 0$)

$$\delta v \sim \exp(i\omega t) = \exp(\Omega t)$$


⇒ strong instability of the radiative driving
(Lucy & Solomon 1970, MacGregor et al. 1979, Carlberg 1980, Owocki et al. 1984)

Wind instabilities III.

- our instability analysis is linear only
- ⇒ hydrodynamical simulations are necessary to describe the instability in detail (Owocki et al. 1988, Feldmeier et al. 1997, Runacres & Owocki 2002)

Wind instabilities III.

- hydrodynamical simulations (Feldmeier et al. 1997)



Wind instabilities III.


- hydrodynamical simulations are able to explain the main properties of X-ray emission of hot stars

Hot star winds: micro-view

- stellar wind of hot stars is accelerated due to the scattering of radiation in lines and on free electrons.
- how does it work on a micro-level?

Hot star winds: micro-view

Typical volume with:
1000 H ions




Hot star winds: micro-view

Typical volume with:
1000 H ions

- radiative acceleration due to the line absorption can be in most cases neglected
- radiative acceleration due to the free-free processes also negligible $\sigma_p \ll \sigma_e$

Hot star winds: micro-view

Typical volume with:
1000 H ions + 100 He ions



Hot star winds: micro-view


Typical volume with:
1000 H ions + 100 He ions

- radiative acceleration due to the line absorption can be in most cases neglected
- radiative acceleration due to the free-free processes also negligible

Hot star winds: micro-view

Typical volume with:

1000 H ions + 100 He ions + 1200 e⁻



Hot star winds: micro-view

Typical volume with:


1000 H ions + 100 He ions + 1200 e⁻

- $\Gamma = g_e / g_{\text{grav}} \approx 0.1$ for many OB stars \Rightarrow significant contribution to the radiative acceleration

Hot star winds: micro-view

Typical volume with:

1000 H ions + 100 He ions + 1200 e⁻ + 2 metals



Hot star winds: micro-view

Typical volume with:


1000 H ions + 100 He ions + 1200 e⁻ + 2 metals

- maximum radiative acceleration due to the lines
 $g_{\text{line}}^{\text{max}} \approx 1000 g_{\text{grav}}$ (Gayley 1995) \Rightarrow crucial contribution to the radiative acceleration

Hot star winds: micro-view

Typical volume with:

1000 H ions + 100 He ions + 1200 e⁻ + 2 metals



How can this work?

two efficient processes necessary:

How can this work?

two efficient processes necessary:

- process which transfers momentum from radiative field to heavier ions

How can this work?

two efficient processes necessary:

- process which transfers momentum from radiative field to heavier ions
- process which transfers momentum from heavier ions to the bulk flow (H, He – mostly passive component)

How to transfer momentum?

- wind is ionised \Rightarrow Coulomb collisions are efficient to transfer momentum from heavier elements to the passive component.

How to transfer momentum?


frictional force on passive component (p) due to ions (i)

$$f_{pi} = \rho_p g_{pi} = n_p n_i \frac{4\pi q_p^2 q_i^2}{k T_{ip}} \ln \Lambda G(x_{ip}) \frac{v_i - v_p}{|v_i - v_p|},$$

where n_p , n_i are number densities of components, v_i , v_p are their radial velocities, and q_p , q_i their charges.

$$x_{ip} = \frac{|v_i - v_p|}{\alpha_{ip}}$$


$$\alpha_{ip}^2 = \frac{2k (m_i T_p + m_p T_i)}{m_i m_p}$$



Momentum transfer efficiency

$$x_{ip} = \frac{|v_{ri} - v_{rp}|}{\alpha_{ip}}$$

$$\alpha_{ip}^2 = \frac{2k (m_i T_p + m_p T_i)}{m_i m_p}$$




- efficient transfer of momentum from heavier ions: one-component models sufficient

Momentum transfer efficiency

$$x_{ip} = \frac{|V_{ri} - V_{rp}|}{\alpha_{ip}}$$

$$\alpha_{ip}^2 = \frac{2k (m_i T_p + m_p T_i)}{m_i m_p}$$




- inefficient transfer of momentum from heavier ions: $x_{ip} \gtrsim 0.1$, part of energy goes to heating – **frictional heating**

Momentum transfer efficiency

$$x_{ip} = \frac{|v_{ri} - v_{rp}|}{\alpha_{ip}}$$

$$\alpha_{ip}^2 = \frac{2k (m_i T_p + m_p T_i)}{m_i m_p}$$



- inefficient collisions between components: $x_{ip} \gtrsim 1$. Chandrasekhar function is a decreasing function of velocity difference \Rightarrow **dynamical decoupling of wind components**
- important for low-density winds (Springmann & Pauldrach 1992, Krtička & Kubát 2001, Votruba et al. 2007).

Hot chemically peculiar stars

- hotter main sequence O stars have winds accelerated by the line transitions of heavier elements (C, N, O, Si, Fe, . . .)

Hot chemically peculiar stars

- for late B stars and A stars (of the main sequence) the radiative force is not strong enough to drive a wind

Hot chemically peculiar stars

- for late B stars and A stars (of the main sequence) the radiative force is not strong enough to drive a wind
- however: the radiative force may cause diffusion of some elements whereas other elements settle down due to the gravity force

Hot chemically peculiar stars

- for late B stars and A stars (of the main sequence) the radiative force is not strong enough to drive a wind
- however: the radiative force may cause diffusion of some elements whereas other elements settle down due to the gravity force

radiative diffusion \times gravitation settling

Hot chemically peculiar stars

- for late B stars and A stars (of the main sequence) the radiative force is not strong enough to drive a wind
- however: the radiative force may cause diffusion of some elements whereas other elements settle down due to the gravity force

radiative diffusion \times gravitation settling

\Rightarrow chemically peculiar (CP) stars

Hot chemically peculiar stars

- for late B stars and A stars (of the main sequence) the radiative force is not strong enough to drive a wind
 - however: the radiative force may cause diffusion of some elements whereas other elements settle down due to the gravity force

radiative diffusion \times gravitation settling
- \Rightarrow chemically peculiar (CP) stars
- overabundance (or underabundance) of certain elements (He, Si, Mg, Fe, ...) in the atmosphere (e.g., Vauclair 2003, Michaud 2005)

Hot chemically peculiar stars

- for late B stars and A stars (of the main sequence) the radiative force is not strong enough to drive a wind
 - however: the radiative force may cause diffusion of some elements whereas other elements settle down due to the gravity force
 - radiative diffusion \times gravitation settling
- \Rightarrow chemically peculiar (CP) stars
- the chemical peculiarity affects surface layers only (the initial chemical composition of the stellar core is roughly solar one)

Hot chemically peculiar stars

- example: HD 37776


Hot chemically peculiar stars

- example: HD 37776




Hot chemically peculiar stars

- example: HD 37776
- Si surface distribution (Chochlova et al. 2000)




Stars in HR diagram




evolutionary tracks (Schearer et al. 1992)

Stars in HR diagram




position of stars discussed here

Stars in HR diagram




stars with P Cyg profiles (Püsküllü et al. 2008)

Stars in HR diagram



stars with different type of wind (Krtićka et al. 2008)

Stars in HR diagram



stars more massive than $M \gtrsim 20 M_{\odot}$ have strong winds basically during all evolutionary phases

The importance of hot star wind I.

- stars more massive than $M \gtrsim 20 M_{\odot}$ have strong winds basically during all evolutionary phases

The importance of hot star wind I.

- stars more massive than $M \gtrsim 20 M_{\odot}$ have strong winds basically during all evolutionary phases
- the duration of the main-sequence phase of massive stars is about 10^6 yr

The importance of hot star wind I.

- stars more massive than $M \gtrsim 20 M_{\odot}$ have strong winds basically during all evolutionary phases
- the duration of the main-sequence phase of massive stars is about 10^6 yr
- during this time massive stars lose mass at the rate of the order of $10^{-6} M_{\odot} \text{ yr}^{-1}$

The importance of hot star wind I.

- stars more massive than $M \gtrsim 20 M_{\odot}$ have strong winds basically during all evolutionary phases
- the duration of the main-sequence phase of massive stars is about 10^6 yr
- during this time massive stars lose mass at the rate of the order of $10^{-6} M_{\odot} \text{ yr}^{-1}$
- a significant part of stellar mass can be lost due to the winds


The importance of hot star wind II

- the evolutionary phases connected with the wind

The importance of hot star wind II

- the evolutionary phases connected with the wind
- Wolf-Rayett stars
 - hot stars with very strong wind (mass-loss rate could be of the order of $10^{-5} M_{\odot} \text{ yr}^{-1}$)
 - wind starts already in the stellar atmosphere
 - spectrum dominated by emission lines
 - enhanced abundance of nitrogen and/or carbon and oxygen

The importance of hot star wind II



The importance of hot star wind II

- the evolutionary phases connected with the wind
- Wolf-Rayett stars
 - how can these stars originate?

The importance of hot star wind II

- the evolutionary phases connected with the wind
- Wolf-Rayett stars
 - during the stellar evolution the core abundance of nitrogen (hydrogen burning) and carbon+oxygen (helium burning) increases

The importance of hot star wind II

- the evolutionary phases connected with the wind
- Wolf-Rayett stars
 - during the stellar evolution the core abundance of nitrogen (hydrogen burning) and carbon+oxygen (helium burning) increases
 - stellar wind blows out the hydrogen-rich stellar envelope and expose nitrogen or carbon+oxygen rich core

The importance of hot star wind II

- the evolutionary phases connected with the wind
 - Wolf-Rayett stars
 - during the stellar evolution the core abundance of nitrogen (hydrogen burning) and carbon+oxygen (helium burning) increases
 - stellar wind blows out the hydrogen-rich stellar envelope and expose nitrogen or carbon+oxygen rich core
- ⇒ Wolf-Rayett stars

The importance of hot star wind II


- the
win

- Wo

-

-

⇒



the

urning)

)

rich

n or

The importance of hot star wind II

- planetary nebulae

The importance of hot star wind II

- planetary nebulae
 - during the AGB stage of solar-like stars ($M \approx 1 M_{\odot}$) the star loses a significant part of its mass via slow ($\sim 10 \text{ km s}^{-1}$) high-density wind

The importance of hot star wind II

- planetary nebulae
 - during the AGB stage of solar-like stars ($M \approx 1 M_{\odot}$) the star loses a significant part of its mass via slow ($\sim 10 \text{ km s}^{-1}$) high-density wind
 - the hot degenerated core is exposed

The importance of hot star wind II

- planetary nebulae
 - during the AGB stage of solar-like stars ($M \approx 1 M_{\odot}$) the star loses a significant part of its mass via slow ($\sim 10 \text{ km s}^{-1}$) high-density wind
 - the hot degenerated core is exposed
 - during this stage the star has fast low-density line-driven wind

The importance of hot star wind II

- planetary nebulae
 - during the AGB stage of solar-like stars ($M \approx 1 M_{\odot}$) the star loses a significant part of its mass via slow ($\sim 10 \text{ km s}^{-1}$) high-density wind
 - the hot degenerated core is exposed
 - during this stage the star has fast low-density line-driven wind
- ⇒ planetary nebula: interaction of slow high-density and fast low-density winds

The importance of hot star wind II

- planetary nebulae



The importance of hot star wind IV

- hot star wind influence also the interstellar environment

(e.g., Dale & Bonnell 2008)

The importance of hot star wind IV

- hot star wind influence also the interstellar environment
 - enrichment of the interstellar medium

(e.g., Dale & Bonnell 2008)


The importance of hot star wind I

- hot star wind influence also the interstellar environment
 - enrichment of the interstellar medium
 - momentum input to the interstellar medium

(e.g., Dale & Bonnell 2008)

The importance of hot star wind IV

-



um
(2008)

What is unclear. . .

- chance for you!

What is unclear. . .

- the most uncertain quantity is . . .

What is unclear. . .

- the most uncertain quantity is the wind mass-loss rate!

What is unclear. . .

- the most uncertain quantity is the wind mass-loss rate!
- why?

What is unclear...

- mass-loss rate and observation

What is unclear. . .

- mass-loss rate and observation
- mass-loss rate can not be derived directly from observation

$$\dot{M} = 4\pi r^2 v \rho$$

- v is fine
- ρ is problematic

What is unclear. . .

- mass-loss rate and observation
- mass-loss rate can not be derived directly from observation
- most of observational characteristics does not depend on ρ , but on ρ^2

What is unclear. . .

- mass-loss rate and observation
- mass-loss rate can not be derived directly from observation
- most of observational characteristics does not depend on ρ , but on ρ^2
- imagine: clumps with the density ρ_c , the mean wind density is $\langle \rho \rangle$

$$\rho_c = C \langle \rho \rangle, \quad C \geq 1$$

What is unclear. . .

- mass-loss rate and observation
- mass-loss rate can not be derived directly from observation
- most of observational characteristics does not depend on ρ , but on ρ^2
- imagine: clumps with the density ρ_c , the mean wind density is $\langle \rho \rangle$

$$\rho_c = C \langle \rho \rangle, \quad C \geq 1$$

$$\langle \rho^2 \rangle = \frac{1}{C} \rho_c^2 = C \langle \rho \rangle^2$$

What is unclear...

- mass-loss rate and observation
- mass-loss rate can not be derived directly from observation
- most of observational characteristics does not depend on ρ , but on ρ^2
- imagine: clumps with the density ρ_c , the mean wind density is $\langle \rho \rangle$

$$\rho_c = C \langle \rho \rangle, \quad C \geq 1$$

$$\langle \rho^2 \rangle_{\text{obs}} = \frac{1}{C} \rho_c^2 = C \langle \rho \rangle_{\text{true}}^2$$

What is unclear...

- mass-loss rate and observation
- mass-loss rate can not be derived directly from observation
- most of observational characteristics does not depend on ρ , but on ρ^2
- imagine: clumps with the density ρ_c , the mean wind density is $\langle \rho \rangle$

$$\rho_c = C \langle \rho \rangle, \quad C \geq 1$$

$$\langle \rho^2 \rangle_{\text{obs}} = \frac{1}{C} \rho_c^2 = C \langle \rho \rangle_{\text{true}}^2$$

⇒ if $C > 1$ we significantly overestimate wind mass-loss rate (by a factor of \sqrt{C})

What is unclear...

- mass-loss rate and theory

What is unclear. . .

- mass-loss rate and theory
- instability of the radiative driving \Rightarrow clumpy wind

What is unclear. . .

- mass-loss rate and theory
- instability of the radiative driving \Rightarrow clumpy wind
- mass-loss rate predicted using smooth wind models


What is unclear. . .

- mass-loss rate and theory
- instability of the radiative driving \Rightarrow clumpy wind
- mass-loss rate predicted using smooth wind models
- what is the influence of inhomogeneities on the predicted mass-loss rates?

What is unclear. . .


- mass-loss rate and theory
 - instability of the radiative driving \Rightarrow clumpy wind
 - mass-loss rate predicted using smooth wind models
 - what is the influence of inhomogeneities on the predicted mass-loss rates?
- \Rightarrow precise values of wind mass-loss rates can not be obtained until we understand the influence of inhomogeneities

What is unclear II.




- what drives winds of WR stars?
(Gräfener & Hamann 2005)

What is unclear III.



- what causes explosions like this?

What is unclear IV.



- what happens outside the well-studied regions?

More informations (papers)

- Milne, E. A. 1926, *MNRAS*, **86**, 459

More informations (papers)

- Milne, E. A. 1926, *MNRAS*, **86**, 459
- Morton, D. C. 1967, *ApJ*, **147**, 1017

More informations (papers)

- Milne, E. A. 1926, *MNRAS*, **86**, 459
- Morton, D. C. 1967, *ApJ*, **147**, 1017
- Lucy, L. B., Solomon, P. M. 1970, *ApJ*, **159**, 879

More informations (papers)

- Milne, E. A. 1926, *MNRAS*, **86**, 459
- Morton, D. C. 1967, *ApJ*, **147**, 1017
- Lucy, L. B., Solomon, P. M. 1970, *ApJ*, **159**, 879
- Castor, J. I. 1974, *MNRAS*, **169**, 279

More informations (papers)

- Milne, E. A. 1926, *MNRAS*, **86**, 459
- Morton, D. C. 1967, *ApJ*, **147**, 1017
- Lucy, L. B., Solomon, P. M. 1970, *ApJ*, **159**, 879
- Castor, J. I. 1974, *MNRAS*, **169**, 279
- Castor, J. I., Abbott, D. C., Klein, R. I. 1975, *ApJ*, **195**, 157

More informations (papers)

- Milne, E. A. 1926, *MNRAS*, **86**, 459
- Morton, D. C. 1967, *ApJ*, **147**, 1017
- Lucy, L. B., Solomon, P. M. 1970, *ApJ*, **159**, 879
- Castor, J. I. 1974, *MNRAS*, **169**, 279
- Castor, J. I., Abbott, D. C., Klein, R. I. 1975, *ApJ*, **195**, 157
- Abbott, D. C. 1980, *ApJ*, **242**, 1183

More informations (papers)

- Milne, E. A. 1926, *MNRAS*, **86**, 459
- Morton, D. C. 1967, *ApJ*, **147**, 1017
- Lucy, L. B., Solomon, P. M. 1970, *ApJ*, **159**, 879
- Castor, J. I. 1974, *MNRAS*, **169**, 279
- Castor, J. I., Abbott, D. C., Klein, R. I. 1975, *ApJ*, **195**, 157
- Abbott, D. C. 1980, *ApJ*, **242**, 1183
- Owocki, S. P., Rybicki, G. B. 1984, *ApJ*, **284**, 337

More informations (papers)

- Milne, E. A. 1926, *MNRAS*, **86**, 459
- Morton, D. C. 1967, *ApJ*, **147**, 1017
- Lucy, L. B., Solomon, P. M. 1970, *ApJ*, **159**, 879
- Castor, J. I. 1974, *MNRAS*, **169**, 279
- Castor, J. I., Abbott, D. C., Klein, R. I. 1975, *ApJ*, **195**, 157
- Abbott, D. C. 1980, *ApJ*, **242**, 1183
- Owocki, S. P., Rybicki, G. B. 1984, *ApJ*, **284**, 337
- Pauldrach A., Puls J., Kudritzki R. P. 1986, *A&A*, **164**, 86

More informations (papers)

- Milne, E. A. 1926, *MNRAS*, **86**, 459
- Morton, D. C. 1967, *ApJ*, **147**, 1017
- Lucy, L. B., Solomon, P. M. 1970, *ApJ*, **159**, 879
- Castor, J. I. 1974, *MNRAS*, **169**, 279
- Castor, J. I., Abbott, D. C., Klein, R. I. 1975, *ApJ*, **195**, 157
- Abbott, D. C. 1980, *ApJ*, **242**, 1183
- Owocki, S. P., Rybicki, G. B. 1984, *ApJ*, **284**, 337
- Pauldrach A., Puls J., Kudritzki R. P. 1986, *A&A*, **164**, 86
- Gayley, K. G., Owocki, S. P. 1994, *ApJ*, **434**, 684

More informations (papers)

- Milne, E. A. 1926, *MNRAS*, **86**, 459
- Morton, D. C. 1967, *ApJ*, **147**, 1017
- Lucy, L. B., Solomon, P. M. 1970, *ApJ*, **159**, 879
- Castor, J. I. 1974, *MNRAS*, **169**, 279
- Castor, J. I., Abbott, D. C., Klein, R. I. 1975, *ApJ*, **195**, 157
- Abbott, D. C. 1980, *ApJ*, **242**, 1183
- Owocki, S. P., Rybicki, G. B. 1984, *ApJ*, **284**, 337
- Pauldrach A., Puls J., Kudritzki R. P. 1986, *A&A*, **164**, 86
- Gayley, K. G., Owocki, S. P. 1994, *ApJ*, **434**, 684
- Feldmeier, A., Puls, J., Pauldrach, A. W. A. 1997, *A&A*, **322**, 878

More informations (book, reviews)

- Lamers, H. J. G. L. M. & Cassinelli, J. P., 1999, Introduction to Stellar Winds (Cambridge: Cambridge Univ. Press)

More informations (book, reviews)

- Lamers, H. J. G. L. M. & Cassinelli, J. P., 1999, Introduction to Stellar Winds (Cambridge: Cambridge Univ. Press)
- Kudritzki, R. P., Puls, J. 2000, *ARA&A*, **38**, 613
(<http://www.usm.uni-muenchen.de/people/puls/Puls.html>)
- Owocki, S. P. 2004, EAS Publications Series, Vol. 13, Evolution of Massive Stars, 163
(<http://www.bartol.udel.edu/~owocki/preprints>)
- Krtička, J., Kubát, J. 2007, Active OB-Stars (San Francisco: ASP Conf. Ser), 153
(<http://arxiv.org/abs/astro-ph/0511443>)
- this lecture
<http://www.physics.muni.cz/~krticka/belehrad.pdf>

Advertisement...

International Conference

Brno, Czech Republic

June 8-12, 2008

Binaries - Key to Comprehension of the Universe

<http://astro.physics.muni.cz/binkey/>

Conclusions

- hot star winds are accelerated by the radiative force due to the line transitions of heavier elements (carbon, nitrogen, silicon, iron, . . .)

Conclusions

- hot star winds are accelerated by the radiative force due to the line transitions of heavier elements
- the most important quantity is the mass loss rate (the amount of mass lost by the star per unit of time)

Conclusions

- hot star winds are accelerated by the radiative force due to the line transitions of heavier elements
- the most important quantity is the mass loss rate
- the mass-loss rate depends mainly on the stellar luminosity (for O stars the mass-loss rate is of the order of $10^{-6} M_{\odot} \text{ yr}^{-1}$)

Conclusions

- hot star winds are accelerated by the radiative force due to the line transitions of heavier elements
- the most important quantity is the mass loss rate
- the mass-loss rate depends mainly on the stellar luminosity
- mass-loss influences the stellar evolution and the circumstellar environment