

Kinematics within the central BLR of AGNs

Wolfram Kollatschny, Göttingen

Platamonas, 2009




University Observatory




Institute for Astrophysics


Scale Sizes of an AGN




Extended radio sources — shown is an FR II source with an edge-brightened structure. The FRIs have lower jet velocities and fade-out to the ends.



The host galaxy. Although shown as an early type galaxy with a smooth profile, it could also be highly irregular with multiple nuclei as a result of merging.




The central kpc star formation disk. This strong far infrared emitting zone might be fed by a bar structure, as seems to be the case for NGC1068.




The narrow-line region comprising small but numerous clouds of the interstellar medium ionized by the central AGN core.


Fig. 9.9 Cartoon of the representative scale sizes of an AGN. How we eventually see the object depends on a number of parameters, the main one being the orientation of the obscuring torus with respect to the observer. (Adapted from Blandford, *Active Galactic Nuclei*, Saas-Fee Advanced Course 20, Springer-Verlag, 1990.)




Broad Line Region Size?




The outer extent of the broad-line region and the deep-walled molecular torus which can provide an effective shield of the central AGN, depending on the relative orientation of the observer.



Inside the molecular torus — the VLBI jet becomes self-absorbed closer in, and the low ionization lines of the BLR, which might be the corona of the accretion disk.



The accretion disk which radiates strongly at UV and optical wavelengths. The high ionization clouds of the BLR are excited by the central continuum radiation field.



The black hole. The Schwarzschild radius for a $10^8 M_{\odot}$ black hole is 2 AU (10^{-5} pc). The spin will introduce twisted magnetic field lines and particle acceleration.

radius:

- $10^{-4} \dots 10^{-1}$ pc
- 1 100 light days

at a dist. of 50 Mpc (Virgo):
spatial resolution

$4 \times 10^{-5} \dots 4 \times 10^{-3}$ ''
(0.04 ... 4. mas)

unresolved

R. Blandford

Broad Line Region Structure?




Fig. 13. A schematic two-component model for the BLR. The high ionization lines are emitted in a spherical system of clouds, and are excited by the direct ultraviolet radiation of the central source. The low ionization lines come mainly from the outer regions of the central disk, where most of the line excitation is due to back-scattered, hard ionizing photons. (After Collin-Souffrin, Perry and Dyson (1987), Collin-Souffrin (1987) and Dumont and Collin-Souffrin (1990))

Two component BLR?

Collin-Souffrin et al., 1990






FIG. 1.—Schematic representation of magnetic accretion disk model for broad emission-line region. The accretion disk is ionized in the inner parts but neutral and probably molecular at large radius. Small dense clouds of molecular gas can be radiatively accelerated away from the surface of the disk and flung centrifugally outward along the magnetic field to attain speeds several times the initial Keplerian velocities. When these clouds are exposed to the full UV photoionizing flux, they are heated to temperatures $T \sim 10^4$ K and produce the emission lines. It is possible that these line photons are subsequently scattered by $\sim 10^4$ K electrons, either within a corona or at the disk.

Radiatively accelerated clouds in hydromagnetic wind?

Emmering, Blandford, Shlosman, 1992

AGN working model

NGC 3783



SMBH $\sim 10^8 M_{\odot} \sim 10^{13} cm$

Broad Line Region ($< 1 pc$)

Narrow Line Region ($\sim 100-1000 pc$)

geometry, kinematics?

Study of Variability:


- Extension, Structure*
- Geometry*
- and Kinematics* of the central
Broad Line Region in AGN

- *Black Hole Mass*

in NGC5548, Mrk926, Mrk110

intermediate, broad-line, narrow-line Sey 1

HST Image of NGC5548



NGC5548

$V = 13.7$


$M_V = -20.7$

$z = 0.017$

$\text{FWHM}(H\beta) = 4400 \text{ km s}^{-1}$

25 x 30 arcsec

High and low state spectra of NGC5548




1989

1993

BLR: Idealized Model


Response of BLR clouds on continuum flashes



BLR stratification


Delay by light travel time effects

BLR: Continuum & integ. $H\beta$ line variability



1989 - 2001

NGC 5548



B. Peterson et al., 2002

$H\beta$ delay ~ 20 light days

1989

BLR size and stratification in NGC5548

peak to peak
var. ampl.

lightcurves (1989)

ACF, CCF

opt. Cont.

0.56

H β


0.29

H α

(0.17)

HeII

0.61



delay

H β

19.



H α

(18.)


HeII

7.

Correlation coeff.




Delay τ [days]




BLR size and stratification in NGC5548

higher ionized lines: - broader line widths
- faster response



Time lag (CCFs centroids) for various emission lines

BLR size and stratification in Mrk110



Mrk110

$V = 15.4$

$M_V = -20.6$


$z = 0.036$

$\text{FWHM}(\text{H}\beta) = 1680 \text{ km s}^{-1}$

tidal arm: 35kpc

HET variability campaign of Mrk110


long-term continuum light curve



1987

2000

Mrk110 spectra taken between 1999 Nov. and 2000 May




9.2m Hobby-Eberly Telescope
at McDonald Observatory
S/N >100

Hobby-Eberly Telescope (HET), McDonald




- Univ. of Texas at Austin
- Pennsylvania State Univ.
- Stanford Univ.
- Göttingen Univ.
- München Univ.

HET variability campaign of Mrk110




Mean spectrum of Mrk110
for 24 epochs from Nov. 1999 through
May 2000



Rms spectrum

- the rms spectrum shows the
variable part of the spectrum

HET variability campaign of Mrk110





Continuum and integrated emission line (Balmer, HeII and HeI) light curves

1999 Nov. - 2000 May

BLR size and structure - HET variab. campaign

Mkn110


CCF functions of HeII, HeI and Balmer line light curves with continuum light curve.



Line	T_{cent} [days]
(1)	(2)
HeII λ 4686	$3.9^{+2.8}_{-0.7}$
HeI λ 4471	$11.1^{+6.0}_{-6.0}$
HeI λ 5016	$14.3^{+7.0}_{-7.0}$
HeI λ 5876	$10.7^{+8.0}_{-6.0}$
H γ	$26.5^{+4.6}_{-4.7}$
H β	$24.2^{+3.7}_{-3.3}$
H α	$32.3^{+4.3}_{-4.9}$


stratification

BLR size and stratification in Mrk110



Normalized rms line profiles in velocity space

The rms spectrum shows the variable part of the spectrum



Mean distances of the line emitting regions from central ionizing source as function of FWHM in rms profiles.

The dotted and dashed lines correspond to virial masses of $.8 - 2.9 \times 10^7 M_\odot$ (from bottom to top).

Central Black Hole Mass in Mrk110


Assumption:
emission line clouds
are gravitationally
bound by central
object

$$M = \frac{fV_{\text{FWHM}}^2 c\tau}{G}$$

$c\tau$ = mean dist. of
line em. clouds

V = vel. disp. of clouds
(from rms line width)


f = factor ($\frac{1}{2}$ - 5.5)
(unknown geometry
and kinematics)



Normalized rms line profiles in velocity space

Line	FWHM(rms) [km s ⁻¹]	τ_{cent} [days]	M [10 ⁷ M _⊙]
(1)	(2)	(3)	(4)
HeIIλ4686	4444. ± 200	3.5 ^{+2.} _{-2.}	2.25 ^{+1.63} _{-0.45}
HeIλ5876	2404. ± 100	10.8 ^{+4.} _{-4.}	1.81 ^{+1.36} _{-0.33}
Hβ	1515. ± 100	23.5 ^{+4.} _{-4.}	1.63 ^{+0.33} _{-0.31}
Hα	1315. ± 100	32.5 ^{+4.} _{-4.}	1.64 ^{+0.33} _{-0.35}

BLR size in Mrk926



Mrk926

$$V = 13.8$$

$$M_V = -23.1$$


$$z = 0.047$$

$$\text{FWHM}(\text{H}\beta) = 9000 \text{ km s}^{-1}$$


compact

HET variability campaign of Mrk926

Continuum light curve 2005




Mrk926 spectra taken between 2005 June and 2005 Dec.




9.2m Hobby-Eberly Telescope at
McDonald Observatory
S/N >100

HET variability campaign of Mrk926





Mean spectrum of Mrk926
for 15 epochs from June 2005 through
Dec. 2005



Mean and Rms spectrum

- the rms spectrum shows the
variable part of the spectrum



HET variability campaign of Mrk926




Normalized mean and rms line profiles in velocity space

- the rms spectrum shows the variable part of the spectrum

Lightcurves of Mrk926



HET var. campaign:
Aug. - Dec. , 2005




mean distance of H β line emitting
region: 0.5 ± 2 light days

Kollatschny et al., 2009 in prep.

BLR size, structure and BH mass in Mrk926

CCF functions of Balmer line light curves with respect to the continuum light curve.



BLR temporally not resolved


-mean distance of $H\beta$ line emitting region: 0.5 ± 1 light day

- $H\alpha$: 1.6 ± 1 light day

v = vel. disp. of clouds (from line width) $\sim 8\,600$ km/s

BH mass $\sim 1. - 3. \cdot 10^7 M_{\odot}$ ($f=1.5$)

Balmer line averaged BLR size in AGN



photoion. theory:

$$r = \left(\frac{Q(\text{H})}{4\pi cn_e} \right)^{1/2} \propto L^{1/2}$$

Q = hydrogen-ionizing photons emitted per sec

★ Mrk926

Relationship between luminosity and broad-line region size


$$R_{\text{BLR}} \sim L^{0.65}$$

But intrinsic scatter due to: BLR density, column density, ionizing spectral energy distribution,?

Kaspi et al. 2004

Central Black Hole Masses in AGN

Black hole mass vs. luminosity for AGN




BH mass for 35 reverberation mapped AGN.


--- : lines of constant mass to luminosity ratio
open circles: NLSy1

Peterson et al., 2004

Line profile segments in Mrk926



H α and H β rms profiles in Mrk926




Velocity-delay map of H α
Grey scale: response of line segments

Fig. 15. The 2-D CCF(τ, v) shows the correlation of the H α line segment light curves with the continuum light curve at 5180 \AA as a function of velocity and time delay (grey scale). Contours of the correlation coefficient are overplotted at levels of 0.85, 0.75, and 0.65 (solid lines). The heavy dashed line connects the centers of all individual cross-correlation functions.


BLR temporally not resolved

Kollatschny et al., 2009 in prep.

Line profile segments in Mrk926




H α and H β rms profiles in Mrk926




Max. response of 2D-CCFs

BLR Kinematics: Idealized Model

Influence of BLR motions on line profile variations



Theory : BLR kinematics - line profile variations



2-d transfer functions

$$L(v, t) = \Psi(v, \tau) * C(t)$$

Ψ now also includes information on the velocity flow

The velocity information is essential: it breaks the degeneracy in the geometry.

theoretical emission line profile variations to derive 2-dim. velocity-delay maps Ψ

velocity-delay maps for different flows


time delay [days]

Welsh & Horne, 1991
Horne et al., 2004

vel. [km/s]

BLR kinematics and accretion disk structure


Mean H β line profile of Mrk110 in velocity space




Light curves of the continuum, of the H β line center, and of different blue and red line wing segments

$$\Delta v = 400 \text{ km/s}$$


Kollatschny & Bischoff 2002



BLR: Accretion disk structure in Mrk110




Mean H β line profile of Mrk110 in velocity space



Velocity-delay map

Kollatschny 2003a

BLR: Accretion disk structure in Mrk110



2-D CCF : correlation of H β line profile segments with cont. variations (grey scale)

Contours of correlation coefficient at levels of .85 to .925 (solid lines).

Dashed curves: theoretical escape velocity envelopes for masses of 0.5, 1., 2. $10^7 M_{\odot}$ (from bottom to top).


Velocity-delay map

Kollatschny 2003a


Theoretical velocity-delay maps for different flows: Keplerian disk BLR model: fast response of **both** outer line wings

Welsh & Horne 1991, Horne et al. 2004

Echo image



Velocity-delay maps: accretion disk structure





2-D CCF : correlation of H β , HeI, HeII line profile segments with continuum variations (grey scale).


Kollatschny 2003

Keplerian disk BLR model: fast response of **both** outer line wings


Solid line: innermost radius at 5 l_d



BLR: Accretion disk wind in Mrk110



2-D CCF: velocity-delay map



Time delay of blue line wing to red line wing as function of dist. to line center

Outer line wings: inner BLR

Disk wind model of BLR: Slightly faster and stronger response of red wing
Chiang & Murray, 1996

Disk driven outflow models compared to spherical wind models: velocity decreases with radius (rather than the other way around)

Koenigl & Kartje, 1994

BLR Structure and Kinematics in Mrk110





FIG. 13.—Schematic representation of how a disk-driven hydromagnetic wind, which is characterized by a highly stratified density distribution, interacts with the active galactic nucleus (AGN) continuum emission. The innermost regions are heated and ionized by the powerful radiation field, with the temperature and degree of ionization varying both with distance and with the polar angle, whereas the outer regions (beyond the dust sublimation radius) are cooler and contain dust. The radiation pressure force on the dust causes the outer streamlines to have a larger opening angle.

Koenigl & Kartje 1994

accretion disk wind in Mrk110

Kollatschny 2003a